

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 November 2001 (29.11.2001)

PCT

(10) International Publication Number
WO 01/90325 A2

- (51) International Patent Classification⁷: C12N 9/00 (US). WILLIAMSON, Mark [US/US]; 15 Stonecrest Drive, Saugus, MA 01906 (US).
- (21) International Application Number: PCT/US01/16549 (74) Agents: MYERS, Louis et al.; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).
- (22) International Filing Date: 21 May 2001 (21.05.2001) (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/205,508 19 May 2000 (19.05.2000) US (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application: US 60/205,508 (CIP) Filed on 19 May 2000 (19.05.2000)
- (71) Applicant (*for all designated States except US*): MILLENNIUM PHARMACEUTICALS, INC. [US/US]; 75 Sidney Street, Cambridge, MA 02139 (US). Published: — *without international search report and to be republished upon receipt of that report*
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): MEYERS, Rachel, A. [US/US]; 115 Devonshire Road, Newton, MA 02468 *For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: 50365, A NOVEL HEXOKINASE FAMILY MEMBER AND USES THEREFOR

(57) Abstract: The invention provides isolated nucleic acids molecules, designated 50365 nucleic acid molecules, which encode novel hexokinase members. The invention also provides antisense nucleic acid molecules, recombinant expression vectors containing 50365 nucleic acid molecules, host cells into which the expression vectors have been introduced, and nonhuman transgenic animals in which a 50365 gene has been introduced or disrupted. The invention still further provides isolated 50365 proteins, fusion proteins, antigenic peptides and anti-50365 antibodies. Diagnostic methods utilizing compositions of the invention are also provided.

WO 01/90325 A2

50365, A NOVEL HEXOKINASE FAMILY MEMBER AND
USES THEREFOR

Related Applications

- 5 [01] This application claims priority to U.S. provisional application number 60/205,508 filed on May 19, 2000, the contents of which are incorporated herein by reference.

Background of the Invention

[02] Hexokinases are a family of sugar phosphorylating enzymes which carry out
10 the phosphorylation of hexoses, for example, glucose, mannose, fructose, sorbitol and glucosamine, at the 6'-position. The phosphoryl donor can be MgATP, ITP, or dATP. Both α - and β -D-hexoses can be phosphorylated, although with different kinetic constants.

[03] Four major isoenzymes are found in vertebrates: types I, II, III, and IV. The liver hexokinase isoenzyme (type IV) is also misleadingly known as glucokinase, and is
15 expressed only in the liver and pancreatic β -cells. This isoenzyme has an important role in modulating insulin secretion. Structurally, the enzymes typically include a small N-terminal hydrophobic region, followed by two similar hexokinase domains of about 450 residues each. The second such region has catalytic activity, while the first has a regulatory role.

[04] Hexokinases are present in nearly all cells. These enzymes have been
20 identified as important for normal glycolytic activity. Irregularities in their function can lead to disorders such as diabetes and hemolytic anemia arising from hexokinase deficiency.

Summary of the Invention

[05] The present invention is based, in part, on the discovery of a novel hexokinase family member, referred to herein as "Fbh50365FL" or "50365". The
25 nucleotide sequence of a cDNA encoding 50365 is shown in SEQ ID NO:1, and the amino acid sequence of a 50365 polypeptide is shown in SEQ ID NO:2 (See also Example 1, below). In addition, the nucleotide sequences of the coding region are recited in SEQ ID NO:3.

[06] Accordingly, in one aspect, the invention features a nucleic acid molecule
30 that encodes a 50365 protein or polypeptide, e.g., a biologically active portion of the 50365

protein. In a preferred embodiment the isolated nucleic acid molecule encodes a polypeptide having the amino acid sequence of SEQ ID NO:2. In other embodiments, the invention provides isolated 50365 nucleic acid molecules having the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, or the sequence of the DNA insert of the plasmid deposited with ATCC Accession Number _____. In still other embodiments, the invention provides nucleic acid molecules that are substantially identical (e.g., naturally occurring allelic variants) to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, or the sequence of the DNA insert of the plasmid deposited with ATCC Accession Number _____. In other embodiments, the invention provides a nucleic acid molecule which hybridizes under a stringency condition described herein to a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, or the sequence of the DNA insert of the plasmid deposited with ATCC Accession Number _____, wherein the nucleic acid encodes a full length 50365 protein or an active fragment thereof.

[07] In a related aspect, the invention further provides nucleic acid constructs that include a 50365 nucleic acid molecule described herein. In certain embodiments, the nucleic acid molecules of the invention are operatively linked to native or heterologous regulatory sequences. Also included, are vectors and host cells containing the 50365 nucleic acid molecules of the invention e.g., vectors and host cells suitable for producing 50365 nucleic acid molecules and polypeptides.

[08] In another related aspect, the invention provides nucleic acid fragments suitable as primers or hybridization probes for the detection of 50365-encoding nucleic acids.

[09] In still another related aspect, isolated nucleic acid molecules that are antisense to a 50365 encoding nucleic acid molecule are provided.

[10] In another aspect, the invention features, 50365 polypeptides, and biologically active or antigenic fragments thereof that are useful, e.g., as reagents or targets in assays applicable to treatment and diagnosis of 50365-mediated or -related disorders. In another embodiment, the invention provides 50365 polypeptides having a 50365 activity. Preferred polypeptides are 50365 proteins including at least one hexokinase domain, and, preferably, having a 50365 activity, e.g., a 50365 activity as described herein.

[11] In other embodiments, the invention provides 50365 polypeptides, e.g., a 50365 polypeptide having the amino acid sequence shown in SEQ ID NO:2 or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with ATCC Accession

Number ____; an amino acid sequence that is substantially identical to the amino acid sequence shown in SEQ ID NO:2 or the amino acid sequence encoded by the cDNA insert of the plasmid deposited with ATCC Accession Number ____; or an amino acid sequence encoded by a nucleic acid molecule having a nucleotide sequence which hybridizes under a stringency condition described herein to a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, or the sequence of the DNA insert of the plasmid deposited with ATCC Accession Number ____, wherein the nucleic acid encodes a full length 50365 protein or an active fragment thereof.

[12] In a related aspect, the invention further provides nucleic acid constructs which include a 50365 nucleic acid molecule described herein.

[13] In a related aspect, the invention provides 50365 polypeptides or fragments operatively linked to non-50365 polypeptides to form fusion proteins.

In another aspect, the invention features antibodies and antigen-binding fragments thereof, that react with, or more preferably specifically bind 50365 polypeptides or fragments thereof, e.g., a hexokinase domain of a 50365 polypeptide. In one embodiment, the antibodies or antigen-binding fragment thereof competitively inhibit the binding of a second antibody to a 50365 polypeptide or fragment thereof, e.g., a hexokinase domain of a 50365 polypeptide.

[14] In another aspect, the invention provides methods of screening for compounds that modulate the expression or activity of the 50365 polypeptides or nucleic acids.

[15] In still another aspect, the invention provides a process for modulating 50365 polypeptide or nucleic acid expression or activity, e.g. using the screened compounds. In certain embodiments, the methods involve treatment of conditions related to aberrant activity or expression of the 50365 polypeptides or nucleic acids, such as conditions involving aberrant or deficient cellular sugar metabolism, e.g., glycolysis.

[16] In yet another aspect, the invention provides methods for inhibiting the proliferation or inducing the killing, of a 50365-expressing cell, e.g., a hyperproliferative 50365-expressing cell. The method includes contacting the cell with an agent, e.g., a compound (e.g., a compound identified using the methods described herein) that modulates the activity, or expression, of the 50365 polypeptide or nucleic acid. In a preferred embodiment, the contacting step is effective *in vitro* or *ex vivo*. In other embodiments, the

contacting step is effected *in vivo*, e.g., in a subject (e.g., a mammal, e.g., a human), as part of a therapeutic or prophylactic protocol.

[17] In a preferred embodiment, the cell is a hyperproliferative cell, e.g., a cell found in a solid tumor, a soft tissue tumor, or a metastatic lesion. In a preferred embodiment, the cell originates from a colon (e.g., a colon tumor or colonic liver metastasis), liver, lung, or ovary cell.

[18] In a preferred embodiment, the agent, e.g., the compound is an inhibitor of a 50365 polypeptide. Preferably, the inhibitor is chosen from a peptide, a phosphopeptide, a small organic molecule, a small inorganic molecule and an antibody (e.g., an antibody conjugated to a therapeutic moiety selected from a cytotoxin, a cytotoxic agent and a radioactive metal ion). In another preferred embodiment, the compound is an inhibitor of a 50365 nucleic acid, e.g., an antisense, a ribozyme, or a triple helix molecule. In still another embodiment, the compound is a substrate analog, e.g., a hexose analog or derivative

[19] In a preferred embodiment, the compound is administered in combination with a cytotoxic agent. Examples of cytotoxic agents include anti-microtubule agent, a topoisomerase I inhibitor, a topoisomerase II inhibitor, an anti-metabolite, a mitotic inhibitor, an alkylating agent, an intercalating agent, an agent capable of interfering with a signal transduction pathway, an agent that promotes apoptosis or necrosis, and radiation.

[20] In another aspect, the invention features methods for treating or preventing a disorder characterized by aberrant cellular proliferation or differentiation of a 50365-expressing cell, in a subject. Preferably, the method includes comprising administering to the subject (e.g., a mammal, e.g., a human) an effective amount of an agent, e.g., a compound (e.g., a compound identified using the methods described herein) that modulates the activity, or expression, of the 50365 polypeptide or nucleic acid. In a preferred embodiment, the disorder is a cancerous or pre-cancerous condition.

[21] In a further aspect, the invention provides methods for evaluating the efficacy of a treatment of a disorder, e.g., proliferative disorder. The method includes: treating a subject, e.g., a patient or an animal, with a protocol under evaluation (e.g., treating a subject with one or more of: chemotherapy, radiation, and/or a compound identified using the methods described herein); and evaluating the expression of a 50365 nucleic acid or polypeptide before and after treatment. A change, e.g., a decrease or increase, in the level of a 50365 nucleic acid (e.g., mRNA) or polypeptide after treatment, relative to the level of expression before treatment, is indicative of the efficacy of the treatment of the disorder.

The level of 50365 nucleic acid or polypeptide expression can be detected by any method described herein.

[22] In a preferred embodiment, the evaluating step includes obtaining a sample (e.g., a tissue sample, e.g., a biopsy, or a fluid sample) from the subject, before and after treatment and comparing the level of expressing of a 50365 nucleic acid (e.g., mRNA) or polypeptide before and after treatment.

[23] In another aspect, the invention provides methods for evaluating the efficacy of a therapeutic or prophylactic agent (e.g., an anti-neoplastic agent). The method includes: contacting a sample with an agent (e.g., a compound identified using the methods described herein, a cytotoxic agent) and, evaluating the expression of 50365 nucleic acid or polypeptide in the sample before and after the contacting step. A change, e.g., a decrease or increase, in the level of 50365 nucleic acid (e.g., mRNA) or polypeptide in the sample obtained after the contacting step, relative to the level of expression in the sample before the contacting step, is indicative of the efficacy of the agent. The level of 50365 nucleic acid or polypeptide expression can be detected by any method described herein. In a preferred embodiment, the sample includes cells obtained from a cancerous tissue or a colon (e.g., normal colon, colon tumor; or colonic liver metastases), liver, lung, or ovary cell tissue.

[24] The invention also provides assays for determining the activity of or the presence or absence of 50365 polypeptides or nucleic acid molecules in a biological sample, including for disease diagnosis.

[25] In further aspect, the invention provides assays for determining the presence or absence of a genetic alteration in a 50365 polypeptide or nucleic acid molecule, including for disease diagnosis.

[26] In one aspect, the invention provides a method of evaluating a sample. The method includes: providing a sample; detecting a 50365 polypeptide or nucleic acid in the sample; and, optionally, comparing the level of expressed 50365 molecules to a reference sample. In one embodiment, an increased level of 50365 molecules is an indication that the sample includes cells in mitosis. In another embodiment, the level of 50365 molecules is an indication that a sample includes a proliferating cell, e.g., a proliferating colon (e.g., normal colon, colon tumor; colonic liver metastases), liver, lung, or ovary cell, preferably a proliferating colon cell.

[27] In another aspect, the invention features a two dimensional array having a plurality of addresses, each address of the plurality being positionally distinguishable from

each other address of the plurality, and each address of the plurality having a unique capture probe, e.g., a nucleic acid or peptide sequence. At least one address of the plurality has a capture probe that recognizes a 50365 molecule. In one embodiment, the capture probe is a nucleic acid, e.g., a probe complementary to a 50365 nucleic acid sequence. In another
5 embodiment, the capture probe is a polypeptide, e.g., an antibody specific for 50365 polypeptides. Also featured is a method of analyzing a sample by contacting the sample to the aforementioned array and detecting binding of the sample to the array.

[28] Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

10 Brief Description of the Drawings

[29] *Figure 1* depicts a hydropathy plot of human 50365. Relative hydrophobic residues are shown above the dashed horizontal line, and relative hydrophilic residues are below the dashed horizontal line. Numbers corresponding to positions in the amino acid sequence of human 50365 are indicated. Polypeptides of the invention include fragments
15 which include: all or part of a hydrophobic sequence, i.e., a sequence above the dashed line, e.g., the sequence of from about amino acid residue 365 to about amino acid residue 380, or from about amino acid residue 645 to about amino acid residue 655, of SEQ ID NO:2; all or part of a hydrophilic sequence, i.e., a sequence below the dashed line, e.g., the sequence of from about amino acid residue 98 to about amino acid residue 120, or from about amino acid
20 residue 715 to about amino acid residue 745 of SEQ ID NO:2. The two hexokinase domains are indicated.

[30] *Figures 2A, 2B, 2C and 2D* depict an alignment of the two hexokinase domains of 50365 with a consensus amino acid sequence derived from a hidden Markov model (PFAM Accession PF00349). The upper sequence is the consensus amino acid
25 sequence (SEQ ID NO:4), while the lower amino acid sequence corresponds to amino acids 16 to 463 (*Figures 2A and 2B*) and amino acids 464 to 910 of SEQ ID NO:2 (*Figures 2C and 2D*).

Detailed Description

[31] The human 50365 sequence (see SEQ ID NO:1, as recited in Example 1),
30 which is approximately 3669 nucleotides long including untranslated regions, contains a

predicted methionine-initiated coding sequence of about 2754 nucleotides, including the termination codon. The coding sequence encodes a 917 amino acid protein (see SEQ ID NO:2, as recited in Example 1).

- [32] Human 50365 contains the following regions or other structural features:
- 5 [33] two hexokinase domains located at about amino acids 16 to 463 and 464 to 910 of SEQ ID NO:2, the latter of which includes a predicted hexokinase signature domain (PFAM Accession Number PS00378) from about amino acid residue 597 to about amino acid residue 622 of SEQ ID NO:2;
- [34] two N-glycosylation sites (PS00001) from about amino acid 208 to about 211, and from about amino acid 655 to about 658, of SEQ ID NO:2;
- 10 [35] one glycosaminoglycan attachment site (PS00002) from about amino acid 896 to about 899 of SEQ ID NO:2;
- [36] one cAMP- and cGMP-dependent protein kinase site (PS00004) from about amino acid 500 to 503 of SEQ ID NO:2;
- 15 [37] twelve protein kinase C phosphorylation sites (PS00005) from about amino acid 172 to 174, 379 to 381, 449 to 451, 508 to 510, 523 to 525, 547 to 549, 551 to 553, 772 to 774, 791 to 793, 826 to 828, 877 to 879, and 896 to 898, of SEQ ID NO:2;
- [38] thirteen casein kinase II sites (PS00006) from about amino acid 35 to 38, 114 to 117, 161 to 164, 243 to 246, 275 to 278, 364 to 367, 569 to 572, 625 to 628, 722 to 725, 20 726 to 729, 787 to 790, 810 to 813, and 877 to 880, of SEQ ID NO:2;
- [39] two tyrosine kinase phosphorylation sites (PS00007) from about amino acid 20 to 27, and 490 to 497, of SEQ ID NO:2;
- [40] twenty-five N-myristylation sites (PS00008) from about amino acid 74 to 79, 151 to 156, 166 to 171, 179 to 184, 212 to 217, 227 to 232, 233 to 238, 299 to 304, 317 to 25 322, 348 to 353, 360 to 365, 411 to 416, 448 to 453, 518 to 523, 589 to 594, 613 to 618, 659 to 664, 674 to 679, 680 to 685, 746 to 751, 779 to 784, 807 to 812, 834 to 839, 858 to 863, and 895 to 900, of SEQ ID NO:2; and
- [41] two amidation sites (PS00009) from amino acid 100 to 103, and amino acid 547 to 550 of SEQ ID NO:2.
- 30 [42] Human 50365 is predicted to be a soluble, cytoplasmic polypeptide.
- [43] For general information regarding PFAM identifiers, PS prefix and PF prefix domain identification numbers, refer to Sonnhammer *et al.* (1997) *Protein* 28:405-420 and <http://www.psc.edu/general/software/packages/pfam/pfam.html>.

[44] A plasmid containing the nucleotide sequence encoding human 50365 (clone "Fbh50365FL") was deposited with American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, VA 20110-2209, on _____ and assigned Accession Number _____. This deposit will be maintained under the terms of the Budapest Treaty on
5 the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. This deposit was made merely as a convenience for those of skill in the art and is not an admission that a deposit is required under 35 U.S.C. §112.

[45] The 50365 protein contains a significant number of structural characteristics in common with members of the hexokinase family. The term "family" when referring to
10 the protein and nucleic acid molecules of the invention means two or more proteins or nucleic acid molecules having a common structural domain or motif and having sufficient amino acid or nucleotide sequence homology as defined herein. Such family members can be naturally or non-naturally occurring and can be from either the same or different species. For example, a family can contain a first protein of human origin as well as other distinct
15 proteins of human origin, or alternatively, can contain homologues of non-human origin, e.g., rat or mouse proteins. Members of a family can also have common functional characteristics.

[46] A 50365 polypeptide can include a "hexokinase domain" or regions homologous with a "hexokinase domain". Type I, II, and III mammalian hexokinase
20 polypeptides typically include two hexokinase domains. Each domain can form a structural unit that includes features of an α/β sandwich. Each domain can include amino acids with regulatory and/or catalytic functions, e.g., including a pocket for ATP and hexose substrates.

[47] As used herein, the term "hexokinase domain" includes an amino acid sequence of about 300 to about 600 amino acid residues in length and having a bit score for
25 the alignment of the sequence to the hexokinase domain (HMM) of at least 300. Preferably, a hexokinase domain includes at least about 350 to about 500 amino acids, more preferably about 400 to about 490 amino acid residues and has a bit score for the alignment of the sequence to the hexokinase domain (HMM) of at least 500, 600, 700, 800 or greater. The hexokinase domain (HMM) has been assigned the PFAM Accession PF00349
30 (<http://genome.wustl.edu/Pfam/html>). An alignment of the hexokinase domains (amino acids 16 to 463 and 464 to 910 of SEQ ID NO:2) of human 50365 with a consensus amino acid sequence derived from a hidden Markov model is depicted in Figure 2.

[48] In a preferred embodiment 50365 polypeptide or protein has a "hexokinase domain" or a region which includes at least about 500 to about 1200, more preferably about 550 to about 1100 or about 600 to about 1000 amino acid residues and has at least about 60%, 70% 80% 90% 95%, 99%, or 100% homology with a "hexokinase domain," e.g., the
5 hexokinase domain of human 50365 (e.g., residues 597 to 622 of SEQ ID NO:2).

[49] Preferably, the hexokinase domain includes a "hexokinase signature domain". This term refers to a protein domain having an amino acid sequence of at least about 15 to about 30 more preferably about 20 to about 30 or about 24 to about 28 amino acid residues which includes the following amino acid sequence: "L-G-F-T-F-S-F-P-C-x-Q-
10 x-S-I-x-x-G-x-L-I-x-W-T-K-G-F" (SEQ ID NO:5). Preferably, a 50365 polypeptide or protein has a "hexokinase signature domain" or a region which includes and has at least about 60%, 70% 80% 90% 95%, 99%, or 100% homology with a "hexokinase signature domain," e.g., the hexokinase signature domain of human 50365 (e.g., residues 597 to 622 of SEQ ID NO:2).

[50] To identify the presence of a "hexokinase" domain in a 50365 protein sequence, and make the determination that a polypeptide or protein of interest has a particular profile, the amino acid sequence of the protein can be searched against the Pfam database of HMMs (e.g., the Pfam database, release 2.1) using the default parameters (http://www.sanger.ac.uk/Software/Pfam/HMM_search). For example, the hmmsf program,
20 which is available as part of the HMMER package of search programs, is a family specific default program for MILPAT0063 and a score of 15 is the default threshold score for determining a hit. Alternatively, the threshold score for determining a hit can be lowered (e.g., to 8 bits). A description of the Pfam database can be found in Sonhammer *et al.* (1997) *Proteins* 28(3):405-420 and a detailed description of HMMs can be found, for
25 example, in Gribskov *et al.* (1990) *Meth. Enzymol.* 183:146-159; Gribskov *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:4355-4358; Krogh *et al.* (1994) *J. Mol. Biol.* 235:1501-1531; and Stultz *et al.* (1993) *Protein Sci.* 2:305-314, the contents of which are incorporated herein by reference. A search was performed against the HMM database resulting in the
30 identification of two "hexokinase domains" in the amino acid sequence of human 50365 at about residues 16 to 463 and 464 to 910 of SEQ ID NO:2.

[51] A 50365 family member can include at least one hexokinase domain. Furthermore, a 50365 family member can include at least one, preferably two N-glycosylation sites (PS00001); at least one glycosaminoglycan attachment site (PS00002); at

least one cAMP- and cGMP-dependent protein kinase site (PS00004); at least one, two, three, four, five, six, seven, eight, nine, ten, eleven, or preferably twelve protein kinase C phosphorylation sites (PS00005); at least one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, or preferably thirteen predicted casein kinase II phosphorylation sites (PS00006); at least one, or preferably two tyrosine kinase phosphorylation sites (PS00007); at least one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, sixteen, seventeen, eighteen, twenty, twenty-three, or preferably twenty-five predicted N-myristylation sites (PS00008); and at least one, preferably two amidation sites (PS00009).

10 [52] As the 50365 polypeptides of the invention may modulate 50365-mediated activities, they may be useful as of for developing novel diagnostic and therapeutic agents for 50365-mediated or related disorders, as described below.

[53] As used herein, a "50365 activity", "biological activity of 50365" or "functional activity of 50365", refers to an activity exerted by a 50365 protein, polypeptide or nucleic acid molecule on e.g., a 50365-responsive cell or on a 50365 substrate, e.g., a protein substrate, as determined *in vivo* or *in vitro*. In one embodiment, a 50365 activity is a direct activity, such as an association with a 50365 target molecule. A "target molecule" or "binding partner" is a molecule with which a 50365 protein binds or interacts in nature. In an exemplary embodiment, is a 50365 substrate, e.g., an aldohexose or ketohexose (e.g., glucose, mannose, fructose, sorbitol and glucosamine), or a phosphate-containing molecule, e.g., ITP, dATP, or MgATP as phosphoryl donor.

20 [54] A 50365 activity can also be an indirect activity, e.g., a cellular signaling activity mediated by interaction of the 50365 protein with a 50365 substrate. For example, the 50365 proteins of the present invention can have one or more of the following activities:

25 (1) it can catalyze the phosphorylation of a sugar, e.g., an aldohexoses and a ketohexoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine); (2) it can catalyze sugar metabolism; (3) it can transfer a phosphate from a phosphate donor (e.g., ATP) to a sugar, e.g., an aldohexoses and a ketohexoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine) to form a phosphorylated sugar, e.g., glucose-6-phosphate; (4) it can modulate glycolytic activities in a cell or tissue, e.g., a tissue in which a 50365 protein is expressed, e.g., muscle tissue and colon; or (5) it can modulate sugar metabolism; and/or (6) it can modulate cellular proliferation and/or differentiation.

30

[55] Based on its structural features, the 50365 molecules of the present invention can have similar biological activities as hexokinase family members.

[56] Expression of 50365 mRNA is modulated in a number of cancerous tissue samples. For example, 50365 mRNA is elevated in a number of colon tumors and colonic
5 liver metastases (see, e.g., Example 2). Thus, the 50365 molecules can act as novel diagnostic targets and therapeutic agents for controlling disorders of neoplasia, e.g., cancer, a cell differentiative disorder, or a cell proliferative disorder. 50365 molecules can also act as indicators and an agent for metabolic disorders, e.g., disorders of sugar metabolism and glycolysis.

10 [57] Examples of cellular proliferative and/or differentiative disorders include cancer, e.g., carcinoma, sarcoma, metastatic disorders or hematopoietic neoplastic disorders, e.g., leukemias. A metastatic tumor can arise from a multitude of primary tumor types, including but not limited to those of prostate, colon, lung, breast and liver origin.

[58] As used herein, the terms "cancer", "hyperproliferative" and "neoplastic"
15 refer to cells having the capacity for autonomous growth. Examples of such cells include cells having an abnormal state or condition characterized by rapidly proliferating cell growth. Hyperproliferative and neoplastic disease states may be categorized as pathologic, i.e., characterizing or constituting a disease state, or may be categorized as non-pathologic, i.e., a deviation from normal but not associated with a disease state. The term is meant to
20 include all types of cancerous growths or oncogenic processes, metastatic tissues or malignantly transformed cells, tissues, or organs, irrespective of histopathologic type or stage of invasiveness. "Pathologic hyperproliferative" cells occur in disease states characterized by malignant tumor growth. Examples of non-pathologic hyperproliferative cells include proliferation of cells associated with wound repair.

25 [59] The terms "cancer" or "neoplasms" include malignancies of the various organ systems, such as affecting lung, breast, thyroid, lymphoid, gastrointestinal, and genitourinary tract, as well as adenocarcinomas which include malignancies such as most colon cancers, renal-cell carcinoma, prostate cancer and/or testicular tumors, non-small cell carcinoma of the lung, cancer of the small intestine and cancer of the esophagus.

30 [60] The term "carcinoma" is art recognized and refers to malignancies of epithelial or endocrine tissues including respiratory system carcinomas, gastrointestinal system carcinomas, genitourinary system carcinomas, testicular carcinomas, breast carcinomas, prostatic carcinomas, endocrine system carcinomas, and melanomas.

Exemplary carcinomas include those forming from tissue of the cervix, lung, prostate, breast, head and neck, colon and ovary. The term also includes carcinosarcomas, e.g., which include malignant tumors composed of carcinomatous and sarcomatous tissues. An “adenocarcinoma” refers to a carcinoma derived from glandular tissue or in which the tumor cells form recognizable glandular structures.

[61] The term “sarcoma” is art recognized and refers to malignant tumors of mesenchymal derivation.

[62] Examples of cellular proliferative and/or differentiative disorders of the colon include, but are not limited to, non-neoplastic polyps, adenomas, familial syndromes, colorectal carcinogenesis, colorectal carcinoma, and carcinoid tumors.

[63] Examples of cellular proliferative and/or differentiative disorders of the liver include, but are not limited to, nodular hyperplasias, adenomas, and malignant tumors, including primary carcinoma of the liver and metastatic tumors.

[64] Examples of cellular proliferative and/or differentiative disorders of the ovary include, but are not limited to, ovarian tumors such as, tumors of coelomic epithelium, serous tumors, mucinous tumors, endometrioid tumors, clear cell adenocarcinoma, cystadenofibroma, brenner tumor, surface epithelial tumors; germ cell tumors such as mature (benign) teratomas, monodermal teratomas, immature malignant teratomas, dysgerminoma, endodermal sinus tumor, choriocarcinoma; sex cord-stromal tumors such as, granulosa-theca cell tumors, thecoma-fibromas, androblastomas, hil cell tumors, and gonadoblastoma; and metastatic tumors such as Krukenberg tumors.

[65] Examples of cellular proliferative and/or differentiative disorders of the lung include, but are not limited to, bronchogenic carcinoma, including paraneoplastic syndromes, bronchioloalveolar carcinoma, neuroendocrine tumors, such as bronchial carcinoid, miscellaneous tumors, and metastatic tumors; pathologies of the pleura, including inflammatory pleural effusions, noninflammatory pleural effusions, pneumothorax, and pleural tumors, including solitary fibrous tumors (pleural fibroma) and malignant mesothelioma.

[66] Examples of cancers or neoplastic conditions, in addition to the ones described above, include, but are not limited to, a fibrosarcoma, myosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing’s tumor, leiomyosarcoma, rhabdomyosarcoma, gastric cancer, esophageal cancer, rectal

cancer, pancreatic cancer, ovarian cancer, prostate cancer, uterine cancer, cancer of the head and neck, skin cancer, brain cancer, squamous cell carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinoma, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma,
5 choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, testicular cancer, small cell lung carcinoma, non-small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, retinoblastoma, leukemia,
10 lymphoma, or Kaposi sarcoma.

[67] Examples of metabolic disorders include insulin resistance like non-insulin-dependent diabetes mellitus (NIDDM), insulin-dependent diabetes mellitus (IDDM), hyperglycaemia, hyperlipidaemia, microvascular angina, congenital muscle fiber type disproportion myopathy, and hemolytic anemia. Metabolic disorders are also further
15 discussed below (see, e.g., "Methods of Treatment").

[68] The 50365 protein, fragments thereof, and derivatives and other variants of the sequence in SEQ ID NO:2 thereof are collectively referred to as "polypeptides or proteins of the invention" or "50365 polypeptides or proteins". Nucleic acid molecules encoding such polypeptides or proteins are collectively referred to as "nucleic acids of the
20 invention" or "50365 nucleic acids." 50365 molecules refer to 50365 nucleic acids, polypeptides, and antibodies.

[69] As used herein, the term "nucleic acid molecule" includes DNA molecules (e.g., a cDNA or genomic DNA), RNA molecules (e.g., an mRNA) and analogs of the DNA or RNA. A DNA or RNA analog can be synthesized from nucleotide analogs. The nucleic
25 acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA.

[70] The term "isolated nucleic acid molecule" or "purified nucleic acid molecule" includes nucleic acid molecules that are separated from other nucleic acid molecules present in the natural source of the nucleic acid. For example, with regards to
30 genomic DNA, the term "isolated" includes nucleic acid molecules which are separated from the chromosome with which the genomic DNA is naturally associated. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and/or 3' ends of the nucleic acid) in the genomic DNA of the

organism from which the nucleic acid is derived. For example, in various embodiments, the isolated nucleic acid molecule can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of 5' and/or 3' nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. Moreover, an
5 "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized.

[71] As used herein, the term "hybridizes under low stringency, medium stringency, high stringency, or very high stringency conditions" describes conditions for
10 hybridization and washing. Guidance for performing hybridization reactions can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6, which is incorporated by reference. Aqueous and nonaqueous methods are described in that reference and either can be used. Specific hybridization conditions referred to herein are as follows: 1) low stringency hybridization conditions in 6X sodium chloride/sodium citrate
15 (SSC) at about 45°C, followed by two washes in 0.2X SSC, 0.1% SDS at least at 50°C (the temperature of the washes can be increased to 55°C for low stringency conditions); 2) medium stringency hybridization conditions in 6X SSC at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 60°C; 3) high stringency hybridization conditions in 6X SSC at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 65°C;
20 and preferably 4) very high stringency hybridization conditions are 0.5M sodium phosphate, 7% SDS at 65°C, followed by one or more washes at 0.2X SSC, 1% SDS at 65°C. Very high stringency conditions (4) are the preferred conditions and the ones that should be used unless otherwise specified.

[72] Preferably, an isolated nucleic acid molecule of the invention that hybridizes
25 under a stringency condition described herein to the sequence of SEQ ID NO:1 or SEQ ID NO:3, corresponds to a naturally-occurring nucleic acid molecule.

[73] As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature. For example a naturally occurring nucleic acid molecule can encode a natural protein.

30 [74] As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules which include at least an open reading frame encoding a 50365 protein. The gene can optionally further include non-coding sequences, e.g., regulatory sequences and introns. Preferably, a gene encodes a mammalian 50365 protein or derivative thereof.

[75] An "isolated" or "purified" polypeptide or protein is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. "Substantially free" means that a preparation of 50365 protein is at least 10% pure. In a preferred embodiment, the preparation of 50365 protein has less than about 30%, 20%, 10% and more preferably 5% (by dry weight), of non-50365 protein (also referred to herein as a "contaminating protein"), or of chemical precursors or non-50365 chemicals. When the 50365 protein or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, i.e., culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation. The invention includes isolated or purified preparations of at least 0.01, 0.1, 1.0, and 10 milligrams in dry weight.

[76] A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence of 50365 without abolishing or substantially altering a 50365 activity. Preferably the alteration does not substantially alter the 50365 activity, e.g., the activity is at least 20%, 40%, 60%, 70% or 80% of wild-type. An "essential" amino acid residue is a residue that, when altered from the wild-type sequence of 50365, results in abolishing a 50365 activity such that less than 20% of the wild-type activity is present. For example, conserved amino acid residues in 50365 are predicted to be particularly unamenable to alteration.

[77] A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in a 50365 protein is preferably replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of a 50365 coding sequence, such as by saturation mutagenesis,

and the resultant mutants can be screened for 50365 biological activity to identify mutants that retain activity. Following mutagenesis of SEQ ID NO:1 or SEQ ID NO:3, the encoded protein can be expressed recombinantly and the activity of the protein can be determined.

[78] As used herein, a "biologically active portion" of a 50365 protein includes a
5 fragment of a 50365 protein which participates in an interaction between a 50365 molecule and a non-50365 molecule. Biologically active portions of a 50365 protein include peptides comprising amino acid sequences sufficiently homologous to or derived from the amino acid sequence of the 50365 protein, e.g., the amino acid sequence shown in SEQ ID NO:2, which include less amino acids than the full length 50365 proteins, and exhibit at least one activity
10 of a 50365 protein. Typically, biologically active portions comprise a domain or motif with at least one activity of the 50365 protein, e.g., catalytic phosphorylation of a sugar molecule, e.g., an aldo- and ketohexoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine); transfer of a phosphate group from a phosphoryl donor, e.g., ITP, dATP or MgATP. A biologically active portion of a 50365 protein can be a polypeptide which is, for example,
15 10, 25, 50, 100, 200 or more amino acids in length. Biologically active portions of a 50365 protein can be used as targets for developing agents which modulate a 50365 mediated activity, e.g., phosphorylation of aldo- and hexoketoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine).

[79] Calculations of homology or sequence identity between sequences (the terms
20 are used interchangeably herein) are performed as follows.

[80] To determine the percent identity of two amino acid sequences, or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for
25 comparison purposes). In a preferred embodiment, the length of a reference sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more preferably at least 50%, 60%, and even more preferably at least 70%, 80%, 90%, 100% of the length of the reference sequence (e.g., when aligning a second sequence to the 50365 amino acid sequence of SEQ ID NO:2 having 918 amino acid residues, at least 40, preferably at least
30 50, more preferably at least 60, even more preferably at least 80, and even more preferably at least 100, or 200 amino acid residues are aligned). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue

or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

5 [81] The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

10 [82] The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch ((1970) *J. Mol. Biol.* 48:444-453) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. A particularly preferred set of parameters (and the one that should be used unless otherwise specified) are a Blossum 62 scoring matrix with a gap penalty of 12, a gap extend penalty of 4, and a frameshift gap penalty of 5.

20 [83] The percent identity between two amino acid or nucleotide sequences can be determined using the algorithm of E. Meyers and W. Miller ((1989) CABIOS, 4:11-17) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

30 [84] The nucleic acid and protein sequences described herein can be used as a "query sequence" to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to 50365 nucleic

acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to 50365 protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) *Nucleic Acids Res.* 25:3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

[85] Particularly preferred 50365 polypeptides of the present invention have an amino acid sequence substantially identical to the amino acid sequence of SEQ ID NO:2. In the context of an amino acid sequence, the term "substantially identical" is used herein to refer to a first amino acid that contains a sufficient or minimum number of amino acid residues that are i) identical to, or ii) conservative substitutions of aligned amino acid residues in a second amino acid sequence such that the first and second amino acid sequences can have a common structural domain and/or common functional activity. For example, amino acid sequences that contain a common structural domain having at least about 60%, or 65% identity, likely 75% identity, more likely 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity to SEQ ID NO:2 are termed substantially identical.

[86] In the context of nucleotide sequence, the term "substantially identical" is used herein to refer to a first nucleic acid sequence that contains a sufficient or minimum number of nucleotides that are identical to aligned nucleotides in a second nucleic acid sequence such that the first and second nucleotide sequences encode a polypeptide having common functional activity, or encode a common structural polypeptide domain or a common functional polypeptide activity. For example, nucleotide sequences having at least about 60%, or 65% identity, likely 75% identity, more likely 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity to SEQ ID NO:1 or 3 are termed substantially identical.

[87] "Misexpression or aberrant expression," as used herein, refers to a non-wildtype pattern of gene expression at the RNA or protein level. It includes: expression at non-wild type levels, i.e., over- or under-expression; a pattern of expression that differs from wild type in terms of the time or stage at which the gene is expressed, e.g., increased or decreased expression (as compared with wild type) at a predetermined developmental period or stage; a pattern of expression that differs from wild type in terms of altered, e.g.,

increased or decreased, expression (as compared with wild type) in a predetermined cell type or tissue type; a pattern of expression that differs from wild type in terms of the splicing size, translated amino acid sequence, post-translational modification, or biological activity of the expressed polypeptide; a pattern of expression that differs from wild type in terms of the effect of an environmental stimulus or extracellular stimulus on expression of the gene, e.g., a pattern of increased or decreased expression (as compared with wild type) in the presence of an increase or decrease in the strength of the stimulus.

[88] "Subject," as used herein, refers to human and non-human animals. The term "non-human animals" of the invention includes all vertebrates, e.g., mammals, such as non-human primates (particularly higher primates), sheep, dog, rodent (e.g., mouse or rat), guinea pig, goat, pig, cat, rabbits, cow, and non-mammals, such as chickens, amphibians, reptiles, etc. In a preferred embodiment, the subject is a human. In another embodiment, the subject is an experimental animal or animal suitable as a disease model.

[89] A "purified preparation of cells," as used herein, refers to an in vitro preparation of cells. In the case cells from multicellular organisms (e.g., plants and animals), a purified preparation of cells is a subset of cells obtained from the organism, not the entire intact organism. In the case of unicellular microorganisms (e.g., cultured cells and microbial cells), it consists of a preparation of at least 10% and more preferably 50% of the subject cells.

[90] Various aspects of the invention are described in further detail below.

Isolated Nucleic Acid Molecules

[91] In one aspect, the invention provides, an isolated or purified, nucleic acid molecule that encodes a 50365 polypeptide described herein, e.g., a full-length 50365 protein or a fragment thereof, e.g., a biologically active portion of 50365 protein. Also included is a nucleic acid fragment suitable for use as a hybridization probe, which can be used, e.g., to identify a nucleic acid molecule encoding a polypeptide of the invention, 50365 mRNA, and fragments suitable for use as primers, e.g., PCR primers for the amplification or mutation of nucleic acid molecules.

[92] In one embodiment, an isolated nucleic acid molecule of the invention includes the nucleotide sequence shown in SEQ ID NO:1, or a portion of any of these nucleotide sequences. In one embodiment, the nucleic acid molecule includes sequences encoding the human 50365 protein (i.e., "the coding region" of SEQ ID NO:1, as shown in

SEQ ID NO:3), as well as 5' untranslated sequences. Alternatively, the nucleic acid molecule can include only the coding region of SEQ ID NO:1 (e.g., nucleotides 107 to 2860, corresponding to SEQ ID NO:3) and, e.g., no flanking sequences which normally accompany the subject sequence. In another embodiment, the nucleic acid molecule encodes a sequence corresponding to the mature protein from about amino acid 1 to amino acid 918 of SEQ ID NO:2.

[93] In another embodiment, an isolated nucleic acid molecule of the invention includes a nucleic acid molecule which is a complement of the nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO:3, or a portion of any of these nucleotide sequences. In other embodiments, the nucleic acid molecule of the invention is sufficiently complementary to the nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO:3, such that it can hybridize (e.g., under a stringency condition described herein) to the nucleotide sequence shown in SEQ ID NO:1 or 3, thereby forming a stable duplex.

[94] In one embodiment, an isolated nucleic acid molecule of the present invention includes a nucleotide sequence which is at least about 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or more homologous to the entire length of the nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO:3, or a portion, preferably of the same length, of any of these nucleotide sequences.

50365 Nucleic Acid Fragments

[95] A nucleic acid molecule of the invention can include only a portion of the nucleic acid sequence of SEQ ID NO:1 or 3. For example, such a nucleic acid molecule can include a fragment which can be used as a probe or primer or a fragment encoding a portion of a 50365 protein, e.g., an immunogenic or biologically active portion of a 50365 protein. A fragment can comprise those nucleotides of SEQ ID NO:1, which encode a hexokinase domain of human 50365. The nucleotide sequence determined from the cloning of the 50365 gene allows for the generation of probes and primers designed for use in identifying and/or cloning other 50365 family members, or fragments thereof, as well as 50365 homologues, or fragments thereof, from other species.

[96] In another embodiment, a nucleic acid includes a nucleotide sequence that includes part, or all, of the coding region and extends into either (or both) the 5' or 3' noncoding region. Other embodiments include a fragment which includes a nucleotide sequence encoding an amino acid fragment described herein. Nucleic acid fragments can encode a specific domain

or site described herein or fragments thereof, particularly fragments thereof which are at least 50, 100, or 250 amino acids in length. Fragments also include nucleic acid sequences corresponding to specific amino acid sequences described above or fragments thereof. Nucleic acid fragments should not to be construed as encompassing those fragments that may have been disclosed prior to the invention.

[97] A nucleic acid fragment can include a sequence corresponding to a domain, region, or functional site described herein. A nucleic acid fragment can also include one or more domain, region, or functional site described herein. Thus, for example, a nucleic acid fragment can comprise nucleotides 1789 to 1866 of SEQ ID NO:3, which encodes a hexokinase signature domain of human 50365.

[98] 50365 probes and primers are provided. Typically a probe/primer is an isolated or purified oligonucleotide. The oligonucleotide typically includes a region of nucleotide sequence that hybridizes under a stringency condition described herein to at least about 7, 12 or 15, preferably about 20 or 25, more preferably about 30, 35, 40, 45, 50, 55, 60, 65, or 75 consecutive nucleotides of a sense or antisense sequence of SEQ ID NO:1 or SEQ ID NO:3, or of a naturally occurring allelic variant or mutant of SEQ ID NO:1 or SEQ ID NO:3.

[99] In a preferred embodiment the nucleic acid is a probe which is at least 5 or 10, and less than 200, more preferably less than 100, or less than 50, base pairs in length. It should be identical, or differ by 1, or less than in 5 or 10 bases, from a sequence disclosed herein. If alignment is needed for this comparison the sequences should be aligned for maximum homology. "Looped" out sequences from deletions or insertions, or mismatches, are considered differences.

[100] A probe or primer can be derived from the sense or anti-sense strand of a nucleic acid which encodes the hexokinase domain, e.g., about nucleotides 106 to 2856 of SEQ ID NO:1 (SEQ ID NO:3), or any other domain, region, or sequence described herein as human 50365.

[101] In another embodiment a set of primers is provided, e.g., primers suitable for use in a PCR, which can be used to amplify a selected region of a 50365 sequence, e.g., a domain, region, site or other sequence described herein. The primers should be at least 5, 10, or 50 base pairs in length and less than 100, or less than 200, base pairs in length. The primers should be identical, or differs by one base from a sequence disclosed herein or from a naturally occurring variant. For example, primers suitable for amplifying all or a portion

of any of the following regions are provided: a hexokinase domain from about amino acid 16 to 463 or 464 to 910 of SEQ ID NO:2.

[102] A nucleic acid fragment can encode an epitope bearing region of a polypeptide described herein.

5 [103] A nucleic acid fragment encoding a "biologically active portion of a 50365 polypeptide" can be prepared by isolating a portion of the nucleotide sequence of SEQ ID NO:1 or 3, which encodes a polypeptide having a 50365 biological activity (e.g., the biological activities of the 50365 proteins are described herein), expressing the encoded portion of the 50365 protein (e.g., by recombinant expression *in vitro*) and assessing the
10 activity of the encoded portion of the 50365 protein. For example, a nucleic acid fragment encoding a biologically active portion of 50365 includes a hexokinase domain, e.g., amino acid residues about 16 to 463 or 464 to 910 of SEQ ID NO:2. A nucleic acid fragment encoding a biologically active portion of a 50365 polypeptide, may comprise a nucleotide sequence which is greater than 300 or more nucleotides in length.

15 [104] In preferred embodiments, a nucleic acid includes a nucleotide sequence which is about 300, 400, 500, 600, 700, 800, 1000, 1200, 1400, 1600, 1800, 2000, 2200, 2400, 2600, 2800, 3000 or more nucleotides in length and hybridizes under a stringency condition described herein to a nucleic acid molecule of SEQ ID NO:1, or SEQ ID NO:3. The nucleic acid can include at least one contiguous nucleotide from a region of about
20 nucleotides 1 to 15, 10 to 700, 666 to 832, 1440 to 2000, 1700 to 2500, 2000 to 2500, 2500 to 3000, 2750 to 3250, or 3000 to 3600 of SEQ ID NO:1.

50365 Nucleic Acid Variants

[105] The invention further encompasses nucleic acid molecules that differ from the nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO:3. Such differences can be
25 due to degeneracy of the genetic code (and result in a nucleic acid which encodes the same 50365 proteins as those encoded by the nucleotide sequence disclosed herein. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence which differs, by at least 1, but less than 5, 10, 20, 50, or 100 amino acid residues that shown in SEQ ID NO:2. If alignment is
30 needed for this comparison the sequences should be aligned for maximum homology. "Looped" out sequences from deletions or insertions, or mismatches, are considered differences.

[106] Nucleic acids of the inventor can be chosen for having codons, which are preferred, or non-preferred, for a particular expression system. E.g., the nucleic acid can be one in which at least one codon, at preferably at least 10%, or 20% of the codons has been altered such that the sequence is optimized for expression in *E. coli*, yeast, human, insect, or
5 CHO cells.

[107] Nucleic acid variants can be naturally occurring, such as allelic variants (same locus), homologs (different locus), and orthologs (different organism) or can be non naturally occurring. Non-naturally occurring variants can be made by mutagenesis techniques, including those applied to polynucleotides, cells, or organisms. The variants can contain nucleotide
10 substitutions, deletions, inversions and insertions. Variation can occur in either or both the coding and non-coding regions. The variations can produce both conservative and non-conservative amino acid substitutions (as compared in the encoded product).

[108] In a preferred embodiment, the nucleic acid differs from that of SEQ ID NO: 1 or 3, e.g., as follows: by at least one but less than 10, 20, 30, or 40 nucleotides; at least one but
15 less than 1%, 5%, 10% or 20% of the nucleotides in the subject nucleic acid. If necessary for this analysis the sequences should be aligned for maximum homology. "Looped" out sequences from deletions or insertions, or mismatches, are considered differences.

[109] Orthologs, homologs, and allelic variants can be identified using methods known in the art. These variants comprise a nucleotide sequence encoding a polypeptide that is
20 50%, at least about 55%, typically at least about 70-75%, more typically at least about 80-85%, and most typically at least about 90-95% or more identical to the nucleotide sequence shown in SEQ ID NO:2 or a fragment of this sequence. Such nucleic acid molecules can readily be identified as being able to hybridize under a stringency condition described herein, to the nucleotide sequence shown in SEQ ID NO 2 or a fragment of the sequence. Nucleic acid
25 molecules corresponding to orthologs, homologs, and allelic variants of the 50365 cDNAs of the invention can further be isolated by mapping to the same chromosome or locus as the 50365 gene.

[110] Preferred variants include those that are correlated with hexokinase activity.

[111] Allelic variants of 50365, e.g., human 50365, include both functional and
30 non-functional proteins. Functional allelic variants are naturally occurring amino acid sequence variants of the 50365 protein within a population that maintain the ability to bind and phosphorylate aldo- and ketohexoses. Functional allelic variants will typically contain only conservative substitution of one or more amino acids of SEQ ID NO:2, or substitution,

deletion or insertion of non-critical residues in non-critical regions of the protein. Non-functional allelic variants are naturally-occurring amino acid sequence variants of the 50365, e.g., human 50365, protein within a population that do not have the ability to bind and phosphorylate a sugar molecule (e.g., an aldo- or a ketohexose, e.g., a glucose, mannose, fructose, sorbitol and glucosamine). Non-functional allelic variants will typically contain a non-conservative substitution, a deletion, or insertion, or premature truncation of the amino acid sequence of SEQ ID NO:2, or a substitution, insertion, or deletion in critical residues or critical regions of the protein.

[112] Moreover, nucleic acid molecules encoding other 50365 family members and, thus, which have a nucleotide sequence which differs from the 50365 sequences of SEQ ID NO:1 or SEQ ID NO:3 are intended to be within the scope of the invention.

Antisense Nucleic Acid Molecules, Ribozymes and Modified 50365 Nucleic Acid Molecules

[113] In another aspect, the invention features, an isolated nucleic acid molecule which is antisense to 50365. An "antisense" nucleic acid can include a nucleotide sequence which is complementary to a "sense" nucleic acid encoding a protein, e.g., complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. The antisense nucleic acid can be complementary to an entire 50365 coding strand, or to only a portion thereof (e.g., the coding region of human 50365 corresponding to SEQ ID NO:3). In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding 50365 (e.g., the 5' and 3' untranslated regions).

[114] An antisense nucleic acid can be designed such that it is complementary to the entire coding region of 50365 mRNA, but more preferably is an oligonucleotide which is antisense to only a portion of the coding or noncoding region of 50365 mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of 50365 mRNA, e.g., between the -10 and +10 regions of the target gene nucleotide sequence of interest. An antisense oligonucleotide can be, for example, about 7, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, or more nucleotides in length.

[115] An antisense nucleic acid of the invention can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example,

an antisense nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used. The antisense nucleic acid also can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

10 [116] The antisense nucleic acid molecules of the invention are typically administered to a subject (e.g., by direct injection at a tissue site), or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a 50365 protein to thereby inhibit expression of the protein, e.g., by inhibiting transcription and/or translation. Alternatively, antisense nucleic acid molecules can be modified to target
15 selected cells and then administered systemically. For systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, e.g., by linking the antisense nucleic acid molecules to peptides or antibodies which bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To
20 achieve sufficient intracellular concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

[117] In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule
25 forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gaultier *et al.* (1987) *Nucleic Acids Res.* 15:6625-6641). The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (Inoue *et al.* (1987) *Nucleic Acids Res.* 15:6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.* (1987) *FEBS Lett.* 215:327-330).

30 [118] In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. A ribozyme having specificity for a 50365-encoding nucleic acid can include one or more sequences complementary to the nucleotide sequence of a 50365 cDNA disclosed herein (i.e., SEQ ID NO:1 or SEQ ID NO:3), and a sequence having known catalytic

sequence responsible for mRNA cleavage (see U.S. Pat. No. 5,093,246 or Haselhoff and Gerlach (1988) *Nature* 334:585-591). For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a 50365-encoding mRNA. See, e.g., Cech *et al.* U.S. Patent No. 4,987,071; and Cech *et al.* U.S. Patent No. 5,116,742. Alternatively, 50365 mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, e.g., Bartel, D. and Szostak, J.W. (1993) *Science* 261:1411-1418.

[119] 50365 gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of the 50365 (e.g., the 50365 promoter and/or enhancers) to form triple helical structures that prevent transcription of the 50365 gene in target cells. See generally, Helene, C. (1991) *Anticancer Drug Des.* 6:569-84; Helene, C. i (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher, L.J. (1992) *Bioassays* 14:807-15. The potential sequences that can be targeted for triple helix formation can be increased by creating a so-called "switchback" nucleic acid molecule. Switchback molecules are synthesized in an alternating 5'-3', 3'-5' manner, such that they base pair with first one strand of a duplex and then the other, eliminating the necessity for a sizeable stretch of either purines or pyrimidines to be present on one strand of a duplex.

[120] The invention also provides detectably labeled oligonucleotide primer and probe molecules. Typically, such labels are chemiluminescent, fluorescent, radioactive, or colorimetric.

[121] A 50365 nucleic acid molecule can be modified at the base moiety, sugar moiety or phosphate backbone to improve, e.g., the stability, hybridization, or solubility of the molecule. For non-limiting examples of synthetic oligonucleotides with modifications see Toulmé (2001) *Nature Biotech.* 19:17 and Faria *et al.* (2001) *Nature Biotech.* 19:40-44. Such phosphoramidite oligonucleotides can be effective antisense agents.

[122] For example, the deoxyribose phosphate backbone of the nucleic acid molecules can be modified to generate peptide nucleic acids (see Hyrup B. *et al.* (1996) *Bioorganic & Medicinal Chemistry* 4: 5-23). As used herein, the terms "peptide nucleic acid" or "PNA" refers to a nucleic acid mimic, e.g., a DNA mimic, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of a PNA can allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of

PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described in Hyrup B. *et al.* (1996) *supra* and Perry-O'Keefe *et al.* *Proc. Natl. Acad. Sci.* 93: 14670-675.

[123] PNAs of 50365 nucleic acid molecules can be used in therapeutic and
5 diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, for example, inducing transcription or translation arrest or inhibiting replication. PNAs of 50365 nucleic acid molecules can also be used in the analysis of single base pair mutations in a gene, (e.g., by PNA-directed PCR clamping); as 'artificial restriction enzymes' when used in combination with other enzymes,
10 (e.g., S1 nucleases (Hyrup B. *et al.* (1996) *supra*)); or as probes or primers for DNA sequencing or hybridization (Hyrup B. *et al.* (1996) *supra*; Perry-O'Keefe *supra*).

[124] In other embodiments, the oligonucleotide may include other appended groups such as peptides (e.g., for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, e.g., Letsinger *et al.* (1989) *Proc. Natl. Acad. Sci.*
15 *USA* 86:6553-6556; Lemaitre *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:648-652; PCT Publication No. W088/09810) or the blood-brain barrier (see, e.g., PCT Publication No. W089/10134). In addition, oligonucleotides can be modified with hybridization-triggered cleavage agents (see, e.g., Krol *et al.* (1988) *Bio-Techniques* 6:958-976) or intercalating agents (see, e.g., Zon (1988) *Pharm. Res.* 5:539-549). To this end, the oligonucleotide may
20 be conjugated to another molecule, (e.g., a peptide, hybridization triggered cross-linking agent, transport agent, or hybridization-triggered cleavage agent).

[125] The invention also includes molecular beacon oligonucleotide primer and probe molecules having at least one region which is complementary to a 50365 nucleic acid of the invention, two complementary regions one having a fluorophore and one a quencher
25 such that the molecular beacon is useful for quantitating the presence of the 50365 nucleic acid of the invention in a sample. Molecular beacon nucleic acids are described, for example, in Lizardi *et al.*, U.S. Patent No. 5,854,033; Nazarenko *et al.*, U.S. Patent No. 5,866,336, and Livak *et al.*, U.S. Patent 5,876,930.

Isolated 50365 Polypeptides

30 [126] In another aspect, the invention features, an isolated 50365 protein, or fragment, e.g., a biologically active portion, for use as immunogens or antigens to raise or test (or more generally to bind) anti-50365 antibodies. 50365 protein can be isolated from

cells or tissue sources using standard protein purification techniques. 50365 protein or fragments thereof can be produced by recombinant DNA techniques or synthesized chemically.

[127] Polypeptides of the invention include those which arise as a result of the existence of multiple genes, alternative transcription events, alternative RNA splicing events, and alternative translational and post-translational events. The polypeptide can be expressed in systems, e.g., cultured cells, which result in substantially the same post-translational modifications present when expressed the polypeptide is expressed in a native cell, or in systems which result in the alteration or omission of post-translational modifications, e.g., glycosylation or cleavage, present when expressed in a native cell.

[128] In a preferred embodiment, a 50365 polypeptide has one or more of the following characteristics:

- [129] (i) it has the ability to modulate cell proliferation and/or metabolism, e.g., sugar metabolism;
- 15 [130] (ii) it has a molecular weight, e.g., a deduced molecular weight, preferably ignoring any contribution of post translational modifications, amino acid composition or other physical characteristic of a 50365 polypeptide, e.g., a polypeptide of SEQ ID NO:2;
- [131] (iii) it has an overall sequence similarity of at least 60%, more preferably at least 70, 80, 90, or 95%, with a polypeptide a of SEQ ID NO:2;
- 20 [132] (iv) it can be found in colon tissue (e.g., normal colon, colon tumor; colonic liver metastases);
- [133] (v) it has a hexokinase domain signature which is preferably about 70%, 80%, 90% or 95% with amino acid residues about 597 to 622 of SEQ ID NO:2;
- [134] (vi) it has one, or preferably two hexokinase domains which are preferably about 70%, 80%, 90% or 95% identical to amino acid residues 16 to 463 or 464 to 910 of SEQ ID NO:2; or
- 25 [135] (vii) it has at least 10, preferably 15, and most preferably 23 of the cysteines found amino acid sequence of the native protein.
- [136] In a preferred embodiment the 50365 protein, or fragment thereof, differs from the corresponding sequence in SEQ ID:2. In one embodiment it differs by at least one but by less than 15, 10 or 5 amino acid residues. In another it differs from the corresponding sequence in SEQ ID NO:2 by at least one residue but less than 20%, 15%, 10% or 5% of the residues in it differ from the corresponding sequence in SEQ ID NO:2. (If this comparison
- 30

requires alignment the sequences should be aligned for maximum homology. "Looped" out sequences from deletions or insertions, or mismatches, are considered differences.) The differences are, preferably, differences or changes at a non essential residue or a conservative substitution. In a preferred embodiment the differences are not in amino acids 16 to 463 or 464 to 910 of SEQ ID NO:2. In another preferred embodiment one or more differences are in amino acids 1 to 463, or 464 to 910, of SEQ ID NO:2.

[137] Other embodiments include a protein that contain one or more changes in amino acid sequence, e.g., a change in an amino acid residue which is not essential for activity. Such 50365 proteins differ in amino acid sequence from SEQ ID NO:2, yet retain biological activity.

[138] In one embodiment, the protein includes an amino acid sequence at least about 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or more homologous to SEQ ID NO:2.

[139] A 50365 protein or fragment is provided which varies from the sequence of SEQ ID NO:2 in regions defined by amino acids about 1 to 463 or 464 to 910, and amino acids 623 to 917 by at least one but by less than 15, 10 or 5 amino acid residues in the protein or fragment but which does not differ from SEQ ID NO:2 in regions defined by amino acids about 1 to 463 or 464 to 910. (If this comparison requires alignment the sequences should be aligned for maximum homology. "Looped" out sequences from deletions or insertions, or mismatches, are considered differences.) In some embodiments the difference is at a non-essential residue or is a conservative substitution, while in others the difference is at an essential residue or is a non-conservative substitution.

[140] In one embodiment, a biologically active portion of a 50365 protein includes one or preferably two hexokinase domains. Moreover, other biologically active portions, in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the functional activities of a native 50365 protein.

[141] In a preferred embodiment, the 50365 protein has an amino acid sequence shown in SEQ ID NO:2. In other embodiments, the 50365 protein is substantially identical to SEQ ID NO:2. In yet another embodiment, the 50365 protein is substantially identical to SEQ ID NO:2 and retains the functional activity of the protein of SEQ ID NO:2, as described in detail in the subsections above.

[142] In another preferred embodiment, the 50365 protein has an amino acid sequence that includes a fragment of SEQ ID NO:2, e.g., about amino acids 16 to 463 or about amino acids 464 to 910 of SEQ ID NO:2.

50365 Chimeric or Fusion Proteins

5 [143] In another aspect, the invention provides 50365 chimeric or fusion proteins. As used herein, a 50365 "chimeric protein" or "fusion protein" includes a 50365 polypeptide linked to a non-50365 polypeptide. A "non-50365 polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein which is not substantially homologous to the 50365 protein, e.g., a protein which is different from the 50365 protein
10 and which is derived from the same or a different organism. The 50365 polypeptide of the fusion protein can correspond to all or a portion e.g., a fragment described herein of a 50365 amino acid sequence. In a preferred embodiment, a 50365 fusion protein includes at least one (or two) biologically active portion of a 50365 protein. The non-50365 polypeptide can be fused to the N-terminus or C-terminus of the 50365 polypeptide.

15 [144] The fusion protein can include a moiety which has a high affinity for a ligand. For example, the fusion protein can be a GST-50365 fusion protein in which the 50365 sequences are fused to the C-terminus of the GST sequences. Such fusion proteins can facilitate the purification of recombinant 50365. Alternatively, the fusion protein can be a 50365 protein containing a heterologous signal sequence at its N-terminus. In certain host
20 cells (e.g., mammalian host cells), expression and/or secretion of 50365 can be increased through use of a heterologous signal sequence.

[145] Fusion proteins can include all or a part of a serum protein, e.g., an IgG constant region, or human serum albumin.

[146] The 50365 fusion proteins of the invention can be incorporated into
25 pharmaceutical compositions and administered to a subject *in vivo*. The 50365 fusion proteins can be used to affect the bioavailability of a 50365 substrate. 50365 fusion proteins may be useful therapeutically for the treatment of disorders caused by, for example, (i) aberrant modification or mutation of a gene encoding a 50365 protein; (ii) mis-regulation of the 50365 gene; and (iii) aberrant post-translational modification of a 50365 protein.

30 [147] Moreover, the 50365-fusion proteins of the invention can be used as immunogens to produce anti-50365 antibodies in a subject, to purify 50365 ligands and in

screening assays to identify molecules which inhibit the interaction of 50365 with a 50365 substrate.

[148] Expression vectors are commercially available that already encode a fusion moiety (e.g., a GST polypeptide). A 50365-encoding nucleic acid can be cloned into such
5 an expression vector such that the fusion moiety is linked in-frame to the 50365 protein.

Variants of 50365 Proteins

[149] In another aspect, the invention also features a variant of a 50365 polypeptide, e.g., which functions as an agonist (mimetics) or as an antagonist. Variants of the 50365 proteins can be generated by mutagenesis, e.g., discrete point mutation, the
10 insertion or deletion of sequences or the truncation of a 50365 protein. An agonist of the 50365 proteins can retain substantially the same, or a subset, of the biological activities of the naturally occurring form of a 50365 protein. An antagonist of a 50365 protein can inhibit one or more of the activities of the naturally occurring form of the 50365 protein by, for example, competitively modulating a 50365-mediated activity of a 50365 protein. Thus,
15 specific biological effects can be elicited by treatment with a variant of limited function. Preferably, treatment of a subject with a variant having a subset of the biological activities of the naturally occurring form of the protein has fewer side effects in a subject relative to treatment with the naturally occurring form of the 50365 protein.

[150] Variants of a 50365 protein can be identified by screening combinatorial
20 libraries of mutants, e.g., truncation mutants, of a 50365 protein for agonist or antagonist activity.

[151] Libraries of fragments e.g., N-terminal, C-terminal, or internal fragments, of a 50365 protein coding sequence can be used to generate a variegated population of fragments for screening and subsequent selection of variants of a 50365 protein. Variants in
25 which a cysteine residues is added or deleted or in which a residue which is glycosylated is added or deleted are particularly preferred.

[152] Methods for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property are known in the art. Such methods are adaptable for rapid screening of
30 the gene libraries generated by combinatorial mutagenesis of 50365 proteins. Recursive ensemble mutagenesis (REM), a new technique which enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify

50365 variants (Arkin and Yourvan (1992) *Proc. Natl. Acad. Sci. USA* 89:7811-7815; Delgrave *et al.* (1993) *Protein Engineering* 6:327-331).

[153] Cell based assays can be exploited to analyze a variegated 50365 library. For example, a library of expression vectors can be transfected into a cell line, e.g., a cell line, which ordinarily responds to 50365 in a substrate-dependent manner. The transfected cells are then contacted with 50365 and the effect of the expression of the mutant on signaling by the 50365 substrate can be detected, e.g., by measuring phosphorylation of aldose- and hexoketoses. Plasmid DNA can then be recovered from the cells which score for inhibition, or alternatively, potentiation of signaling by the 50365 substrate, and the individual clones further characterized.

[154] In another aspect, the invention features a method of making a 50365 polypeptide, e.g., a peptide having a non-wild type activity, e.g., an antagonist, agonist, or super agonist of a naturally occurring 50365 polypeptide, e.g., a naturally occurring 50365 polypeptide. The method includes: altering the sequence of a 50365 polypeptide, e.g., altering the sequence, e.g., by substitution or deletion of one or more residues of a non-conserved region, a domain or residue disclosed herein, and testing the altered polypeptide for the desired activity.

[155] In another aspect, the invention features a method of making a fragment or analog of a 50365 polypeptide a biological activity of a naturally occurring 50365 polypeptide. The method includes: altering the sequence, e.g., by substitution or deletion of one or more residues, of a 50365 polypeptide, e.g., altering the sequence of a non-conserved region, or a domain or residue described herein, and testing the altered polypeptide for the desired activity.

Anti-50365 Antibodies

[156] In another aspect, the invention provides an anti-50365 antibody, or a fragment thereof (e.g., an antigen-binding fragment thereof). The term "antibody" as used herein refers to an immunoglobulin molecule or immunologically active portion thereof, i.e., an antigen-binding portion. As used herein, the term "antibody" refers to a protein comprising at least one, and preferably two, heavy (H) chain variable regions (abbreviated herein as VH), and at least one and preferably two light (L) chain variable regions (abbreviated herein as VL). The VH and VL regions can be further subdivided into regions of hypervariability, termed "complementarity determining regions" ("CDR"), interspersed

with regions that are more conserved, termed "framework regions" (FR). The extent of the framework region and CDR's has been precisely defined (see, Kabat, E.A., *et al.* (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242, and Chothia, C. *et al.* (1987) *J. Mol. Biol.* 196:901-917, which are incorporated herein by reference). Each VH and VL is composed of three CDR's and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4.

[157] The anti-50365 antibody can further include a heavy and light chain constant region, to thereby form a heavy and light immunoglobulin chain, respectively. In one embodiment, the antibody is a tetramer of two heavy immunoglobulin chains and two light immunoglobulin chains, wherein the heavy and light immunoglobulin chains are interconnected by, e.g., disulfide bonds. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. The light chain constant region is comprised of one domain, CL. The variable region of the heavy and light chains contains a binding domain that interacts with an antigen. The constant regions of the antibodies typically mediate the binding of the antibody to host tissues or factors, including various cells of the immune system (e.g., effector cells) and the first component (C1q) of the classical complement system.

[158] As used herein, the term "immunoglobulin" refers to a protein consisting of one or more polypeptides substantially encoded by immunoglobulin genes. The recognized human immunoglobulin genes include the kappa, lambda, alpha (IgA1 and IgA2), gamma (IgG1, IgG2, IgG3, IgG4), delta, epsilon and mu constant region genes, as well as the myriad immunoglobulin variable region genes. Full-length immunoglobulin "light chains" (about 25 KDa or 214 amino acids) are encoded by a variable region gene at the NH₂-terminus (about 110 amino acids) and a kappa or lambda constant region gene at the COOH--terminus. Full-length immunoglobulin "heavy chains" (about 50 KDa or 446 amino acids), are similarly encoded by a variable region gene (about 116 amino acids) and one of the other aforementioned constant region genes, e.g., gamma (encoding about 330 amino acids).

[159] The term "antigen-binding fragment" of an antibody (or simply "antibody portion," or "fragment"), as used herein, refers to one or more fragments of a full-length antibody that retain the ability to specifically bind to the antigen, e.g., 50365 polypeptide or fragment thereof. Examples of antigen-binding fragments of the anti-50365 antibody include, but are not limited to: (i) a Fab fragment, a monovalent fragment consisting of the

- VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (v) a dAb fragment (Ward *et al.*, (1989) *Nature* 341:544-546), which consists of a VH domain; and (vi) an isolated complementarity determining region (CDR). Furthermore, although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the VL and VH regions pair to form monovalent molecules (known as single chain Fv (scFv); see *e.g.*, Bird *et al.* (1988) *Science* 242:423-426; and Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883). Such single chain antibodies are also encompassed within the term "antigen-binding fragment" of an antibody. These antibody fragments are obtained using conventional techniques known to those with skill in the art, and the fragments are screened for utility in the same manner as are intact antibodies.
- 15 [160] The anti-50365 antibody can be a polyclonal or a monoclonal antibody. In other embodiments, the antibody can be recombinantly produced, *e.g.*, produced by phage display or by combinatorial methods.
- [161] Phage display and combinatorial methods for generating anti-50365 antibodies are known in the art (as described in, *e.g.*, Ladner *et al.* U.S. Patent No. 5,223,409; Kang *et al.* International Publication No. WO 92/18619; Dower *et al.* International Publication No. WO 91/17271; Winter *et al.* International Publication WO 92/20791; Markland *et al.* International Publication No. WO 92/15679; Breitling *et al.* International Publication WO 93/01288; McCafferty *et al.* International Publication No. WO 92/01047; Garrard *et al.* International Publication No. WO 92/09690; Ladner *et al.* International Publication No. WO 90/02809; Fuchs *et al.* (1991) *Bio/Technology* 9:1370-1372; Hay *et al.* (1992) *Hum Antibod Hybridomas* 3:81-85; Huse *et al.* (1989) *Science* 246:1275-1281; Griffiths *et al.* (1993) *EMBO J* 12:725-734; Hawkins *et al.* (1992) *J Mol Biol* 226:889-896; Clackson *et al.* (1991) *Nature* 352:624-628; Gram *et al.* (1992) *PNAS* 89:3576-3580; Garrad *et al.* (1991) *Bio/Technology* 9:1373-1377; Hoogenboom *et al.* (1991) *Nuc Acid Res* 19:4133-4137; and Barbas *et al.* (1991) *PNAS* 88:7978-7982, the contents of all of which are incorporated by reference herein).
- 30 [162] In one embodiment, the anti-50365 antibody is a fully human antibody (*e.g.*, an antibody made in a mouse which has been genetically engineered to produce an antibody

from a human immunoglobulin sequence), or a non-human antibody, e.g., a rodent (mouse or rat), goat, primate (e.g., monkey), camel antibody. Preferably, the non-human antibody is a rodent (mouse or rat antibody). Method of producing rodent antibodies are known in the art.

- 5 [163] Human monoclonal antibodies can be generated using transgenic mice carrying the human immunoglobulin genes rather than the mouse system. Splenocytes from these transgenic mice immunized with the antigen of interest are used to produce hybridomas that secrete human mAbs with specific affinities for epitopes from a human protein (see, e.g., Wood et al. International Application WO 91/00906, Kucherlapati et al.
- 10 PCT publication WO 91/10741; Lonberg et al. International Application WO 92/03918; Kay et al. International Application 92/03917; Lonberg, N. et al. 1994 *Nature* 368:856-859; Green, L.L. et al. 1994 *Nature Genet.* 7:13-21; Morrison, S.L. et al. 1994 *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Bruggeman et al. 1993 *Year Immunol* 7:33-40; Tuailon et al. 1993 *PNAS* 90:3720-3724; Bruggeman et al. 1991 *Eur J Immunol* 21:1323-1326).
- 15 [164] An anti-50365 antibody can be one in which the variable region, or a portion thereof, e.g., the CDR's, are generated in a non-human organism, e.g., a rat or mouse. Chimeric, CDR-grafted, and humanized antibodies are within the invention. Antibodies generated in a non-human organism, e.g., a rat or mouse, and then modified, e.g., in the variable framework or constant region, to decrease antigenicity in a human are within the
- 20 invention.
- [165] Chimeric antibodies can be produced by recombinant DNA techniques known in the art. For example, a gene encoding the Fc constant region of a murine (or other species) monoclonal antibody molecule is digested with restriction enzymes to remove the region encoding the murine Fc, and the equivalent portion of a gene encoding a human Fc
- 25 constant region is substituted (see Robinson et al., International Patent Publication PCT/US86/02269; Akira, et al., European Patent Application 184,187; Taniguchi, M., European Patent Application 171,496; Morrison et al., European Patent Application 173,494; Neuberger et al., International Application WO 86/01533; Cabilly et al. U.S. Patent No. 4,816,567; Cabilly et al., European Patent Application 125,023; Better et al. (1988
- 30 *Science* 240:1041-1043); Liu et al. (1987) *PNAS* 84:3439-3443; Liu et al., 1987, *J. Immunol.* 139:3521-3526; Sun et al. (1987) *PNAS* 84:214-218; Nishimura et al., 1987, *Canc. Res.* 47:999-1005; Wood et al. (1985) *Nature* 314:446-449; and Shaw et al., 1988, *J. Natl Cancer Inst.* 80:1553-1559).

[166] A humanized or CDR-grafted antibody will have at least one or two but generally all three recipient CDR's (of heavy and or light immunoglobulin chains) replaced with a donor CDR. The antibody may be replaced with at least a portion of a non-human CDR or only some of the CDR's may be replaced with non-human CDR's. It is only
5 necessary to replace the number of CDR's required for binding of the humanized antibody to a 50365 or a fragment thereof. Preferably, the donor will be a rodent antibody, e.g., a rat or mouse antibody, and the recipient will be a human framework or a human consensus framework. Typically, the immunoglobulin providing the CDR's is called the "donor" and the immunoglobulin providing the framework is called the "acceptor." In one embodiment,
10 the donor immunoglobulin is a non-human (e.g., rodent). The acceptor framework is a naturally-occurring (e.g., a human) framework or a consensus framework, or a sequence about 85% or higher, preferably 90%, 95%, 99% or higher identical thereto.

[167] As used herein, the term "consensus sequence" refers to the sequence formed from the most frequently occurring amino acids (or nucleotides) in a family of related
15 sequences (See e.g., Winnaker, *From Genes to Clones* (Verlagsgesellschaft, Weinheim, Germany 1987). In a family of proteins, each position in the consensus sequence is occupied by the amino acid occurring most frequently at that position in the family. If two amino acids occur equally frequently, either can be included in the consensus sequence. A "consensus framework" refers to the framework region in the consensus immunoglobulin sequence.

[168] An antibody can be humanized by methods known in the art. Humanized antibodies can be generated by replacing sequences of the Fv variable region which are not directly involved in antigen binding with equivalent sequences from human Fv variable regions. General methods for generating humanized antibodies are provided by Morrison, S. L., 1985, *Science* 229:1202-1207, by Oi et al., 1986, *BioTechniques* 4:214, and by Queen et
25 al. US 5,585,089, US 5,693,761 and US 5,693,762, the contents of all of which are hereby incorporated by reference. Those methods include isolating, manipulating, and expressing the nucleic acid sequences that encode all or part of immunoglobulin Fv variable regions from at least one of a heavy or light chain. Sources of such nucleic acid are well known to those skilled in the art and, for example, may be obtained from a hybridoma producing an
30 antibody against a 50365 polypeptide or fragment thereof. The recombinant DNA encoding the humanized antibody, or fragment thereof, can then be cloned into an appropriate expression vector.

- [169] Humanized or CDR-grafted antibodies can be produced by CDR-grafting or CDR substitution, wherein one, two, or all CDR's of an immunoglobulin chain can be replaced. See e.g., U.S. Patent 5,225,539; Jones et al. 1986 *Nature* 321:552-525; Verhoeyan et al. 1988 *Science* 239:1534; Beidler et al. 1988 *J. Immunol.* 141:4053-4060; Winter US 5,225,539, the contents of all of which are hereby expressly incorporated by reference. Winter describes a CDR-grafting method which may be used to prepare the humanized antibodies of the present invention (UK Patent Application GB 2188638A, filed on March 26, 1987; Winter US 5,225,539), the contents of which is expressly incorporated by reference.
- 10 [170] Also within the scope of the invention are humanized antibodies in which specific amino acids have been substituted, deleted or added. Preferred humanized antibodies have amino acid substitutions in the framework region, such as to improve binding to the antigen. For example, a humanized antibody will have framework residues identical to the donor framework residue or to another amino acid other than the recipient
- 15 framework residue. To generate such antibodies, a selected, small number of acceptor framework residues of the humanized immunoglobulin chain can be replaced by the corresponding donor amino acids. Preferred locations of the substitutions include amino acid residues adjacent to the CDR, or which are capable of interacting with a CDR (see e.g., US 5,585,089). Criteria for selecting amino acids from the donor are described in US
- 20 5,585,089, e.g., columns 12-16 of US 5,585,089, the e.g., columns 12-16 of US 5,585,089, the contents of which are hereby incorporated by reference. Other techniques for humanizing antibodies are described in Padlan et al. EP 519596 A1, published on December 23, 1992.
- [171] In preferred embodiments an antibody can be made by immunizing with
- 25 purified 50365 antigen, or a fragment thereof, e.g., a fragment described herein.
- [172] A full-length 50365 protein or, antigenic peptide fragment of 50365 can be used as an immunogen or can be used to identify anti-50365 antibodies made with other immunogens, e.g., cells, membrane preparations, and the like. The antigenic peptide of 50365 should include at least 8 amino acid residues of the amino acid sequence shown in
- 30 SEQ ID NO:2 and encompasses an epitope of 50365. Preferably, the antigenic peptide includes at least 10 amino acid residues, more preferably at least 15 amino acid residues, even more preferably at least 20 amino acid residues, and most preferably at least 30 amino acid residues.

[173] Fragments of 50365 that include residues 98 to 120, or 715 to 745 can be used to make, e.g., used as immunogens or used to characterize the specificity of an antibody, antibodies against hydrophilic regions of the 50365 protein. Similarly, a fragment of 50365 which includes residues 16 to 463 or 464 to 910 of SEQ ID NO:2 can be used to make an antibody against a hydrophobic region of the 50365 protein; a fragment of 50365 which includes residues 597 to 622 can be used to make an antibody against the hexokinase signature region of the 50365 protein.

[174] Antibodies reactive with, or specific for, any of these regions, or other regions or domains described herein are provided.

10 [175] Antibodies which bind only native 50365 protein, only denatured or otherwise non-native 50365 protein, or which bind both, are within the invention. Antibodies with linear or conformational epitopes are within the invention. Conformational epitopes can sometimes be identified by identifying antibodies which bind to native but not denatured 50365 protein.

15 [176] Preferred epitopes encompassed by the antigenic peptide are regions of 50365 are located on the surface of the protein, e.g., hydrophilic regions, as well as regions with high antigenicity. For example, an Emini surface probability analysis of the human 50365 protein sequence can be used to indicate the regions that have a particularly high probability of being localized to the surface of the 50365 protein and are thus likely to constitute surface residues useful for targeting antibody production.

20 [177] In a preferred embodiment the antibody binds an epitope on any domain or region on 50365 proteins described herein.

[178] The anti-50365 antibody can be a single chain antibody. A single-chain antibody (scFV) may be engineered (see, for example, Colcher, D. *et al.* (1999) *Ann N Y Acad Sci* 880:263-80; and Reiter, Y. (1996) *Clin Cancer Res* 2:245-52). The single chain antibody can be dimerized or multimerized to generate multivalent antibodies having specificities for different epitopes of the same target 50365 protein.

25 [179] In a preferred embodiment the antibody has: effector function; and can fix complement. In other embodiments the antibody does not; recruit effector cells; or fix complement.

30 [180] In a preferred embodiment, the antibody has reduced or no ability to bind an Fc receptor. For example, it is an isotype or subtype, fragment or other mutant, which does

not support binding to an Fc receptor, e.g., it has a mutagenized or deleted Fc receptor binding region.

[181] In a preferred embodiment, an anti-50365 antibody alters (e.g., increases or decreases) the catalytic phosphorylation of a sugar molecule, e.g., an aldo- and ketohexoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine); and transfer of a phosphate group from a phosphoryl donor, e.g., ITP, dATP or MgATP, activity of a 50365 polypeptide. For example, an anti-50365 antibody can bind to the active site, e.g., at or near a catalytic residue of the 50635 protein.

[182] The antibody can be coupled to a toxin, e.g., a polypeptide toxin, e.g., ricin or diphtheria toxin or active fragment hereof, or a radioactive nucleus, or imaging agent, e.g. a radioactive, enzymatic, or other, e.g., imaging agent, e.g., a NMR contrast agent. Labels which produce detectable radioactive emissions or fluorescence are preferred.

[183] An anti-50365 antibody (e.g., monoclonal antibody) can be used to isolate 50365 by standard techniques, such as affinity chromatography or immunoprecipitation. Moreover, an anti-50365 antibody can be used to detect 50365 protein (e.g., in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the protein. Anti-50365 antibodies can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, e.g., to determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance (i.e., antibody labelling). Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

[184] The invention also includes nucleic acids which encode an anti-50365 antibody, e.g., an anti-50365 antibody described herein. Also included are vectors which include the nucleic acid and cells transformed with the nucleic acid, particularly cells which are useful for producing an antibody, e.g., mammalian cells, e.g. CHO or lymphatic cells.

[185] The invention also includes cell lines, e.g., hybridomas, which make an anti-50365 antibody, e.g., and antibody described herein, and method of using said cells to make a 50365 antibody.

Recombinant Expression Vectors, Host Cells and Genetically Engineered Cells

5 [186] In another aspect, the invention includes, vectors, preferably expression vectors, containing a nucleic acid encoding a polypeptide described herein. As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked and can include a plasmid, cosmid or viral vector. The vector can be capable of autonomous replication or it can integrate into a host DNA. Viral
10 vectors include, e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses.

[187] A vector can include a 50365 nucleic acid in a form suitable for expression of the nucleic acid in a host cell. Preferably the recombinant expression vector includes one or more regulatory sequences operatively linked to the nucleic acid sequence to be expressed.
15 The term "regulatory sequence" includes promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence, as well as tissue-specific regulatory and/or inducible sequences. The design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, and the
20 like. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or polypeptides, including fusion proteins or polypeptides, encoded by nucleic acids as described herein (e.g., 50365 proteins, mutant forms of 50365 proteins, fusion proteins, and the like).

[188] The recombinant expression vectors of the invention can be designed for
25 expression of 50365 proteins in prokaryotic or eukaryotic cells. For example, polypeptides of the invention can be expressed in *E. coli*, insect cells (e.g., using baculovirus expression vectors), yeast cells or mammalian cells. Suitable host cells are discussed further in Goeddel; (1990) *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA. Alternatively, the recombinant expression vector can be transcribed
30 and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

[189] Expression of proteins in prokaryotes is most often carried out in *E. coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion
5 vectors typically serve three purposes: 1) to increase expression of recombinant protein; 2) to increase the solubility of the recombinant protein; and 3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to
10 purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith, D.B. and Johnson, K.S. (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively,
15 to the target recombinant protein.

[190] Purified fusion proteins can be used in 50365 activity assays, (e.g., direct assays or competitive assays described in detail below), or to generate antibodies specific for 50365 proteins. In a preferred embodiment, a fusion protein expressed in a retroviral expression vector of the present invention can be used to infect bone marrow cells which are
20 subsequently transplanted into irradiated recipients. The pathology of the subject recipient is then examined after sufficient time has passed (e.g., six weeks).

[191] To maximize recombinant protein expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., (1990) *Gene Expression Technology: Methods in Enzymology* 185,
25 Academic Press, San Diego, California 119-128). Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in *E. coli* (Wada *et al.*, (1992) *Nucleic Acids Res.* 20:2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

[192] The 50365 expression vector can be a yeast expression vector, a vector for
30 expression in insect cells, e.g., a baculovirus expression vector or a vector suitable for expression in mammalian cells.

[193] When used in mammalian cells, the expression vector's control functions can be provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40.

[194] In another embodiment, the promoter is an inducible promoter, e.g., a promoter regulated by a steroid hormone, by a polypeptide hormone (e.g., by means of a signal transduction pathway), or by a heterologous polypeptide (e.g., the tetracycline-inducible systems, "Tet-On" and "Tet-Off"; see, e.g., Clontech Inc., CA, Gossen and Bujard (1992) *Proc. Natl. Acad. Sci. USA* 89:5547, and Paillard (1989) *Human Gene Therapy* 9:983).

10 [195] In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (e.g., tissue-specific regulatory elements are used to express the nucleic acid). Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert *et al.* (1987) *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and
15 Eaton (1988) *Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore (1989) *EMBO J.* 8:729-733) and immunoglobulins (Banerji *et al.* (1983) *Cell* 33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters (e.g., the neurofilament promoter; Byrne and Ruddle (1989) *Proc. Natl. Acad. Sci. USA* 86:5473-5477), pancreas-specific promoters (Edlund *et al.* (1985) *Science*
20 230:912-916), and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Patent No. 4,873,316 and European Application Publication No. 264,166).

Developmentally-regulated promoters are also encompassed, for example, the murine hox promoters (Kessel and Gruss (1990) *Science* 249:374-379) and the α -fetoprotein promoter (Campes and Tilghman (1989) *Genes Dev.* 3:537-546).

25 [196] The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. Regulatory sequences (e.g., viral promoters and/or enhancers) operatively linked to a nucleic acid cloned in the antisense orientation can be chosen which direct the constitutive, tissue specific or cell type specific expression of antisense RNA in a variety of cell types. The
30 antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus.

[197] Another aspect the invention provides a host cell which includes a nucleic acid molecule described herein, e.g., a 50365 nucleic acid molecule within a recombinant

expression vector or a 50365 nucleic acid molecule containing sequences which allow it to homologously recombine into a specific site of the host cell's genome. The terms "host cell" and "recombinant host cell" are used interchangeably herein. Such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because
5 certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

[198] A host cell can be any prokaryotic or eukaryotic cell. For example, a 50365 protein can be expressed in bacterial cells (such as *E. coli*), insect cells, yeast or mammalian
10 cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

[199] Vector DNA can be introduced into host cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing
15 foreign nucleic acid (e.g., DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation.

[200] A host cell of the invention can be used to produce (i.e., express) a 50365 protein. Accordingly, the invention further provides methods for producing a 50365 protein
20 using the host cells of the invention. In one embodiment, the method includes culturing the host cell of the invention (into which a recombinant expression vector encoding a 50365 protein has been introduced) in a suitable medium such that a 50365 protein is produced. In another embodiment, the method further includes isolating a 50365 protein from the medium or the host cell.

25 [201] In another aspect, the invention features, a cell or purified preparation of cells which include a 50365 transgene, or which otherwise misexpress 50365. The cell preparation can consist of human or non-human cells, e.g., rodent cells, e.g., mouse or rat cells, rabbit cells, or pig cells. In preferred embodiments, the cell or cells include a 50365 transgene, e.g., a heterologous form of a 50365, e.g., a gene derived from humans (in the
30 case of a non-human cell). The 50365 transgene can be misexpressed, e.g., overexpressed or underexpressed. In other preferred embodiments, the cell or cells include a gene that mis-expresses an endogenous 50365, e.g., a gene the expression of which is disrupted, e.g., a

knockout. Such cells can serve as a model for studying disorders that are related to mutated or mis-expressed 50365 alleles or for use in drug screening.

[202] In another aspect, the invention features, a human cell, e.g., a hematopoietic stem cell or a fibroblast cell, transformed with nucleic acid which encodes a subject 50365 polypeptide.

[203] Also provided are cells, preferably human cells, e.g., human hematopoietic or fibroblast cells, in which an endogenous 50365 is under the control of a regulatory sequence that does not normally control the expression of the endogenous 50365 gene. The expression characteristics of an endogenous gene within a cell, e.g., a cell line or microorganism, can be modified by inserting a heterologous DNA regulatory element into the genome of the cell such that the inserted regulatory element is operably linked to the endogenous 50365 gene. For example, an endogenous 50365 gene which is "transcriptionally silent," e.g., not normally expressed, or expressed only at very low levels, may be activated by inserting a regulatory element which is capable of promoting the expression of a normally expressed gene product in that cell. Techniques such as targeted homologous recombinations, can be used to insert the heterologous DNA as described in, e.g., Chappel, US 5,272,071; WO 91/06667, published in May 16, 1991.

[204] In a preferred embodiment, recombinant cells described herein can be used for replacement therapy in a subject. For example, a nucleic acid encoding a 50365 polypeptide operably linked to an inducible promoter (e.g., a steroid hormone receptor-regulated promoter) is introduced into a human or nonhuman, e.g., mammalian, e.g., porcine recombinant cell. The cell is cultivated and encapsulated in a biocompatible material, such as poly-lysine alginate, and subsequently implanted into the subject. See, e.g., Lanza (1996) *Nat. Biotechnol.* 14:1107; Joki *et al.* (2001) *Nat. Biotechnol.* 19:35; and U.S. Patent No. 5,876,742. Production of 50365 polypeptide can be regulated in the subject by administering an agent (e.g., a steroid hormone) to the subject. In another preferred embodiment, the implanted recombinant cells express and secrete an antibody specific for a 50365 polypeptide. The antibody can be any antibody or any antibody derivative described herein.

30 Transgenic Animals

[205] The invention provides non-human transgenic animals. Such animals are useful for studying the function and/or activity of a 50365 protein and for identifying and/or

evaluating modulators of 50365 activity. As used herein, a "transgenic animal" is a non-human animal, preferably a mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, amphibians, and the like. A transgene is exogenous DNA or a rearrangement, e.g., a deletion of endogenous chromosomal DNA, which preferably is integrated into or occurs in the genome of the cells of a transgenic animal. A transgene can direct the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal, other transgenes, e.g., a knockout, reduce expression. Thus, a transgenic animal can be one in which an endogenous 50365 gene has been altered by, e.g., by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, e.g., an embryonic cell of the animal, prior to development of the animal.

[206] Intronic sequences and polyadenylation signals can also be included in the transgene to increase the efficiency of expression of the transgene. A tissue-specific regulatory sequence(s) can be operably linked to a transgene of the invention to direct expression of a 50365 protein to particular cells. A transgenic founder animal can be identified based upon the presence of a 50365 transgene in its genome and/or expression of 50365 mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene encoding a 50365 protein can further be bred to other transgenic animals carrying other transgenes.

[207] 50365 proteins or polypeptides can be expressed in transgenic animals or plants, e.g., a nucleic acid encoding the protein or polypeptide can be introduced into the genome of an animal. In preferred embodiments the nucleic acid is placed under the control of a tissue specific promoter, e.g., a milk or egg specific promoter, and recovered from the milk or eggs produced by the animal. Suitable animals are mice, pigs, cows, goats, and sheep.

[208] The invention also includes a population of cells from a transgenic animal, as discussed, e.g., below.

30 Uses

[209] The nucleic acid molecules, proteins, protein homologues, and antibodies described herein can be used in one or more of the following methods: a) screening assays;

b) predictive medicine (e.g., diagnostic assays, prognostic assays, monitoring clinical trials, and pharmacogenetics); c) methods of treatment (e.g., therapeutic and prophylactic); and d) synthetic methods for producing carbohydrates, e.g., sugars, e.g., hexoses and variants thereof.

- 5 [210] The isolated nucleic acid molecules of the invention can be used, for example, to express a 50365 protein (e.g., via a recombinant expression vector in a host cell in gene therapy applications), to detect a 50365 mRNA (e.g., in a biological sample) or a genetic alteration in a 50365 gene, and to modulate 50365 activity, as described further below. The 50365 proteins can be used to treat disorders characterized by insufficient or
10 excessive production of a 50365 substrate or production of 50365 inhibitors. In addition, the 50365 proteins can be used to screen for naturally occurring 50365 substrates, to screen for drugs or compounds which modulate 50365 activity, as well as to treat disorders characterized by insufficient or excessive production of 50365 protein or production of 50365 protein forms which have decreased, aberrant or unwanted activity compared to
15 50365 wild type protein (e.g., faulty glycolytic activity, including faulty phosphorylation of aldo- and ketohexoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine)). Moreover, the anti-50365 antibodies of the invention can be used to detect and isolate 50365 proteins, regulate the bioavailability of 50365 proteins, and modulate 50365 activity.
- [211] A method of evaluating a compound for the ability to interact with, e.g., bind,
20 a subject 50365 polypeptide is provided. The method includes: contacting the compound with the subject 50365 polypeptide; and evaluating ability of the compound to interact with, e.g., to bind or form a complex with the subject 50365 polypeptide. This method can be performed in vitro, e.g., in a cell free system, or in vivo, e.g., in a two-hybrid interaction trap assay. This method can be used to identify naturally occurring molecules that interact with
25 subject 50365 polypeptide. It can also be used to find natural or synthetic inhibitors of subject 50365 polypeptide. Screening methods are discussed in more detail below.

Screening Assays

- [212] The invention provides methods (also referred to herein as “screening assays”) for identifying modulators, i.e., candidate or test compounds or agents (e.g.,
30 proteins, peptides, peptidomimetics, peptoids, small molecules or other drugs) which bind to 50365 proteins, have a stimulatory or inhibitory effect on, for example, 50365 expression or 50365 activity, or have a stimulatory or inhibitory effect on, for example, the expression or

activity of a 50365 substrate. Compounds thus identified can be used to modulate the activity of target gene products (e.g., 50365 genes) in a therapeutic protocol, to elaborate the biological function of the target gene product, or to identify compounds that disrupt normal target gene interactions.

5 [213] In one embodiment, the invention provides assays for screening candidate or test compounds which are substrates of a 50365 protein or polypeptide or a biologically active portion thereof. In another embodiment, the invention provides assays for screening candidate or test compounds that bind to or modulate an activity of a 50365 protein or polypeptide or a biologically active portion thereof.

10 [214] Any routine method for assaying hexokinase activity can be used to assay the hexokinase activity of a 50365 protein. For example, Liu *et al.* (1999) *J Biol Chem* 274:31155 describe assays of the enzymological properties of a human hexokinase.

[215] The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including:
15 biological libraries; peptoid libraries (libraries of molecules having the functionalities of peptides, but with a novel, non-peptide backbone which are resistant to enzymatic degradation but which nevertheless remain bioactive; see, e.g., Zuckermann, R.N. *et al.* (1994) *J. Med. Chem.* 37:2678-85); spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-
20 compound' library method; and synthetic library methods using affinity chromatography selection. The biological library and peptoid library approaches are limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam (1997) *Anticancer Drug Des.* 12:145).

[216] Examples of methods for the synthesis of molecular libraries can be found in
25 the art, for example in: DeWitt *et al.* (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90:6909; Erb *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:11422; Zuckermann *et al.* (1994) *J. Med. Chem.* 37:2678; Cho *et al.* (1993) *Science* 261:1303; Carrell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2059; Carrell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2061; and Gallop *et al.* (1994) *J. Med. Chem.* 37:1233.

30 [217] Libraries of compounds may be presented in solution (e.g., Houghten (1992) *Biotechniques* 13:412-421), or on beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner, U.S. Patent No. 5,223,409), spores (Ladner U.S. Patent No. 5,223,409), plasmids (Cull *et al.* (1992) *Proc Natl Acad Sci USA* 89:1865-

1869) or on phage (Scott and Smith (1990) *Science* 249:386-390; Devlin (1990) *Science* 249:404-406; Cwirla *et al.* (1990) *Proc. Natl. Acad. Sci.* 87:6378-6382; Felici (1991) *J. Mol. Biol.* 222:301-310; Ladner *supra.*).

[218] In one embodiment, an assay is a cell-based assay in which a cell which
5 expresses a 50365 protein or biologically active portion thereof is contacted with a test compound, and the ability of the test compound to modulate 50365 activity is determined. Determining the ability of the test compound to modulate 50365 activity can be accomplished by monitoring, for example, catalytic phosphorylation of a sugar molecule, e.g., an aldo- and ketohexoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine);
10 transfer of a phosphate group from a phosphoryl donor, e.g., ITP, dATP or MgATP. The cell, for example, can be of mammalian origin, e.g., human.

[219] The ability of the test compound to modulate 50365 binding to a compound, e.g., a 50365 substrate, or to bind to 50365 can also be evaluated. This can be accomplished, for example, by coupling the compound, e.g., the substrate, with a
15 radioisotope or enzymatic label such that binding of the compound, e.g., the substrate, to 50365 can be determined by detecting the labeled compound, e.g., substrate, in a complex. Alternatively, 50365 could be coupled with a radioisotope or enzymatic label to monitor the ability of a test compound to modulate 50365 binding to a 50365 substrate in a complex. For example, compounds (e.g., 50365 substrates) can be labeled with ^{125}I , ^{35}S , ^{14}C , or ^3H ,
20 either directly or indirectly, and the radioisotope detected by direct counting of radioemmission or by scintillation counting. Alternatively, compounds can be enzymatically labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product.

[220] The ability of a compound (e.g., a 50365 substrate) to interact with 50365
25 with or without the labeling of any of the interactants can be evaluated. For example, a microphysiometer can be used to detect the interaction of a compound with 50365 without the labeling of either the compound or the 50365 (McConnell, H. M. *et al.* (1992) *Science* 257:1906-1912). As used herein, a "microphysiometer" (e.g., Cytosensor) is an analytical
30 instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a compound and 50365.

[221] In yet another embodiment, a cell-free assay is provided in which a 50365 protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to bind to the 50365 protein or biologically active portion thereof is evaluated. Preferred biologically active portions of the 50365 proteins to be used
5 in assays of the present invention include fragments which participate in interactions with non-50365 molecules, e.g., fragments with high surface probability scores.

[222] Soluble and/or membrane-bound forms of isolated proteins (e.g., 50365 proteins or biologically active portions thereof) can be used in the cell-free assays of the invention. When membrane-bound forms of the protein are used, it may be desirable to
10 utilize a solubilizing agent. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton® X-100, Triton® X-114, Thesit®, Isotridecypoly(ethylene glycol ether)_n, 3-[(3-cholamidopropyl)dimethylamminio]-1-propane sulfonate (CHAPS), 3-[(3-cholamidopropyl)dimethylamminio]-2-hydroxy-1-propane
15 sulfonate (CHAPSO), or N-dodecyl=N,N-dimethyl-3-ammonio-1-propane sulfonate.

[223] Cell-free assays involve preparing a reaction mixture of the target gene protein and the test compound under conditions and for a time sufficient to allow the two components to interact and bind, thus forming a complex that can be removed and/or detected.

20 [224] The interaction between two molecules can also be detected, e.g., using fluorescence energy transfer (FET) (see, for example, Lakowicz *et al.*, U.S. Patent No. 5,631,169; Stavrianopoulos, *et al.*, U.S. Patent No. 4,868,103). A fluorophore label on the first, 'donor' molecule is selected such that its emitted fluorescent energy will be absorbed by a fluorescent label on a second, 'acceptor' molecule, which in turn is able to fluoresce
25 due to the absorbed energy. Alternately, the 'donor' protein molecule may simply utilize the natural fluorescent energy of tryptophan residues. Labels are chosen that emit different wavelengths of light, such that the 'acceptor' molecule label may be differentiated from that of the 'donor'. Since the efficiency of energy transfer between the labels is related to the distance separating the molecules, the spatial relationship between the molecules can be
30 assessed. In a situation in which binding occurs between the molecules, the fluorescent emission of the 'acceptor' molecule label in the assay should be maximal. An FET binding event can be conveniently measured through standard fluorometric detection means well known in the art (e.g., using a fluorimeter).

- [225] In another embodiment, determining the ability of the 50365 protein to bind to a target molecule can be accomplished using real-time Biomolecular Interaction Analysis (BIA) (see, e.g., Sjolander, S. and Urbaniczky, C. (1991) *Anal. Chem.* 63:2338-2345 and Szabo *et al.* (1995) *Curr. Opin. Struct. Biol.* 5:699-705). "Surface plasmon resonance" or
- 5 "BIA" detects biospecific interactions in real time, without labeling any of the interactants (e.g., BIAcore). Changes in the mass at the binding surface (indicative of a binding event) result in alterations of the refractive index of light near the surface (the optical phenomenon of surface plasmon resonance (SPR)), resulting in a detectable signal which can be used as an indication of real-time reactions between biological molecules.
- 10 [226] In one embodiment, the target gene product or the test substance is anchored onto a solid phase. The target gene product/test compound complexes anchored on the solid phase can be detected at the end of the reaction. Preferably, the target gene product can be anchored onto a solid surface, and the test compound, (which is not anchored), can be labeled, either directly or indirectly, with detectable labels discussed herein.
- 15 [227] It may be desirable to immobilize either 50365, an anti-50365 antibody or its target molecule to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to a 50365 protein, or interaction of a 50365 protein with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable
- 20 for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase/50365 fusion proteins or glutathione-S-transferase/target fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO)
- 25 or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and either the non-adsorbed target protein or 50365 protein, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components, the matrix immobilized in the
- 30 case of beads, complex determined either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of 50365 binding or activity determined using standard techniques.

[228] Other techniques for immobilizing either a 50365 protein or a target molecule on matrices include using conjugation of biotin and streptavidin. Biotinylated 50365 protein or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques known in the art (e.g., biotinylation kit, Pierce Chemicals, Rockford, IL), and
5 immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical).

[229] In order to conduct the assay, the non-immobilized component is added to the coated surface containing the anchored component. After the reaction is complete, unreacted components are removed (e.g., by washing) under conditions such that any complexes formed will remain immobilized on the solid surface. The detection of
10 complexes anchored on the solid surface can be accomplished in a number of ways. Where the previously non-immobilized component is pre-labeled, the detection of label immobilized on the surface indicates that complexes were formed. Where the previously non-immobilized component is not pre-labeled, an indirect label can be used to detect complexes anchored on the surface; e.g., using a labeled antibody specific for the
15 immobilized component (the antibody, in turn, can be directly labeled or indirectly labeled with, e.g., a labeled anti-Ig antibody).

[230] In one embodiment, this assay is performed utilizing antibodies reactive with 50365 protein or target molecules but which do not interfere with binding of the 50365 protein to its target molecule. Such antibodies can be derivatized to the wells of the plate, and unbound target or 50365 protein trapped in the wells by antibody conjugation. Methods
20 for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the 50365 protein or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the 50365 protein or target molecule.

[231] Alternatively, cell free assays can be conducted in a liquid phase. In such an assay, the reaction products are separated from unreacted components, by any of a number of standard techniques, including but not limited to: differential centrifugation (see, for example, Rivas, G., and Minton, A.P., (1993) *Trends Biochem Sci* 18:284-7);
chromatography (gel filtration chromatography, ion-exchange chromatography);
25 electrophoresis (see, e.g., Ausubel, F. *et al.*, eds. *Current Protocols in Molecular Biology* 1999, J. Wiley: New York.); and immunoprecipitation (see, for example, Ausubel, F. *et al.*, eds. (1999) *Current Protocols in Molecular Biology*, J. Wiley: New York). Such resins and chromatographic techniques are known to one skilled in the art (see, e.g., Heegaard, N.H.,
30

(1998) *J Mol Recognit* 11:141-8; Hage, D.S., and Tweed, S.A. (1997) *J Chromatogr B Biomed Sci Appl.* 699:499-525). Further, fluorescence energy transfer may also be conveniently utilized, as described herein, to detect binding without further purification of the complex from solution.

5 [232] In a preferred embodiment, the assay includes contacting the 50365 protein or biologically active portion thereof with a known compound which binds 50365 to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with a 50365 protein, wherein determining the ability of the test compound to interact with a 50365 protein includes determining the ability of the
10 test compound to preferentially bind to 50365 or biologically active portion thereof, or to modulate the activity of a target molecule, as compared to the known compound.

[233] The target gene products of the invention can, *in vivo*, interact with one or more cellular or extracellular macromolecules, such as proteins. For the purposes of this discussion, such cellular and extracellular macromolecules are referred to herein as "binding
15 partners." Compounds that disrupt such interactions can be useful in regulating the activity of the target gene product. Such compounds can include, but are not limited to molecules such as antibodies, peptides, and small molecules. The preferred target genes/products for use in this embodiment are the 50365 genes herein identified. In an alternative embodiment, the invention provides methods for determining the ability of the test compound to modulate
20 the activity of a 50365 protein through modulation of the activity of a downstream effector of a 50365 target molecule. For example, the activity of the effector molecule on an appropriate target can be determined, or the binding of the effector to an appropriate target can be determined, as previously described.

[234] To identify compounds that interfere with the interaction between the target
25 gene product and its cellular or extracellular binding partner(s), a reaction mixture containing the target gene product and the binding partner is prepared, under conditions and for a time sufficient, to allow the two products to form complex. In order to test an inhibitory agent, the reaction mixture is provided in the presence and absence of the test compound. The test compound can be initially included in the reaction mixture, or can be
30 added at a time subsequent to the addition of the target gene and its cellular or extracellular binding partner. Control reaction mixtures are incubated without the test compound or with a placebo. The formation of any complexes between the target gene product and the cellular or extracellular binding partner is then detected. The formation of a complex in the control

reaction, but not in the reaction mixture containing the test compound, indicates that the compound interferes with the interaction of the target gene product and the interactive binding partner. Additionally, complex formation within reaction mixtures containing the test compound and normal target gene product can also be compared to complex formation within reaction mixtures containing the test compound and mutant target gene product. This comparison can be important in those cases wherein it is desirable to identify compounds that disrupt interactions of mutant but not normal target gene products.

[235] These assays can be conducted in a heterogeneous or homogeneous format. Heterogeneous assays involve anchoring either the target gene product or the binding partner onto a solid phase, and detecting complexes anchored on the solid phase at the end of the reaction. In homogeneous assays, the entire reaction is carried out in a liquid phase. In either approach, the order of addition of reactants can be varied to obtain different information about the compounds being tested. For example, test compounds that interfere with the interaction between the target gene products and the binding partners, e.g., by competition, can be identified by conducting the reaction in the presence of the test substance. Alternatively, test compounds that disrupt preformed complexes, e.g., compounds with higher binding constants that displace one of the components from the complex, can be tested by adding the test compound to the reaction mixture after complexes have been formed. The various formats are briefly described below.

[236] In a heterogeneous assay system, either the target gene product or the interactive cellular or extracellular binding partner is anchored onto a solid surface (e.g., a microtiter plate), while the non-anchored species is labeled, either directly or indirectly. The anchored species can be immobilized by non-covalent or covalent attachments. Alternatively, an immobilized antibody specific for the species to be anchored can be used to anchor the species to the solid surface.

[237] In order to conduct the assay, the partner of the immobilized species is exposed to the coated surface with or without the test compound. After the reaction is complete, unreacted components are removed (e.g., by washing) and any complexes formed will remain immobilized on the solid surface. Where the non-immobilized species is pre-labeled, the detection of label immobilized on the surface indicates that complexes were formed. Where the non-immobilized species is not pre-labeled, an indirect label can be used to detect complexes anchored on the surface; e.g., using a labeled antibody specific for the initially non-immobilized species (the antibody, in turn, can be directly labeled or indirectly

labeled with, e.g., a labeled anti-Ig antibody). Depending upon the order of addition of reaction components, test compounds that inhibit complex formation or that disrupt preformed complexes can be detected.

[238] Alternatively, the reaction can be conducted in a liquid phase in the presence
5 or absence of the test compound, the reaction products separated from unreacted components, and complexes detected; e.g., using an immobilized antibody specific for one of the binding components to anchor any complexes formed in solution, and a labeled antibody specific for the other partner to detect anchored complexes. Again, depending upon the order of addition of reactants to the liquid phase, test compounds that inhibit
10 complex or that disrupt preformed complexes can be identified.

[239] In an alternate embodiment of the invention, a homogeneous assay can be used. For example, a preformed complex of the target gene product and the interactive cellular or extracellular binding partner product is prepared in that either the target gene products or their binding partners are labeled, but the signal generated by the label is
15 quenched due to complex formation (see, e.g., U.S. Patent No. 4,109,496 that utilizes this approach for immunoassays). The addition of a test substance that competes with and displaces one of the species from the preformed complex will result in the generation of a signal above background. In this way, test substances that disrupt target gene product-binding partner interaction can be identified.

[240] In yet another aspect, the 50365 proteins can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Patent No. 5,283,317; Zervos *et al.* (1993) *Cell* 72:223-232; Madura *et al.* (1993) *J. Biol. Chem.* 268:12046-12054; Bartel *et al.* (1993) *Biotechniques* 14:920-924; Iwabuchi *et al.* (1993) *Oncogene* 8:1693-1696; and Brent WO94/10300), to identify other proteins, which bind to or interact with 50365
25 ("50365-binding proteins" or "50365-bp") and are involved in 50365 activity. Such 50365-bps can be activators or inhibitors of signals by the 50365 proteins or 50365 targets as, for example, downstream elements of a 50365-mediated signaling pathway.

[241] The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay
30 utilizes two different DNA constructs. In one construct, the gene that codes for a 50365 protein is fused to a gene encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for

the activation domain of the known transcription factor. (Alternatively the: 50365 protein can be the fused to the activator domain.) If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a 50365-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., lacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the 50365 protein.

10 [242] In another embodiment, modulators of 50365 expression are identified. For example, a cell or cell free mixture is contacted with a candidate compound and the expression of 50365 mRNA or protein evaluated relative to the level of expression of 50365 mRNA or protein in the absence of the candidate compound. When expression of 50365 mRNA or protein is greater in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of 50365 mRNA or protein expression. Alternatively, when expression of 50365 mRNA or protein is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of 50365 mRNA or protein expression. The level of 50365 mRNA or protein expression can be determined by methods described herein for detecting 50365 mRNA or protein.

[243] In another aspect, the invention pertains to a combination of two or more of the assays described herein. For example, a modulating agent can be identified using a cell-based or a cell free assay, and the ability of the agent to modulate the activity of a 50365 protein can be confirmed *in vivo*, e.g., in an animal such as an animal model for faulty glycolytic activity, including faulty phosphorylation of aldo- and ketohexoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine).

25 [244] This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein (e.g., a 50365 modulating agent, an antisense 50365 nucleic acid molecule, a 50365-specific antibody, or a 50365-binding partner) in an appropriate animal model to determine the efficacy, toxicity, side effects, or mechanism of action, of treatment with such an agent. Furthermore, novel agents identified by the above-described screening assays can be used for treatments as described herein.

Detection Assays

[245] Portions or fragments of the nucleic acid sequences identified herein can be used as polynucleotide reagents. For example, these sequences can be used to: (i) map their respective genes on a chromosome e.g., to locate gene regions associated with genetic disease or to associate 50365 with a disease; (ii) identify an individual from a minute biological sample (tissue typing); and (iii) aid in forensic identification of a biological sample. These applications are described in the subsections below.

Chromosome Mapping

[246] The 50365 nucleotide sequences or portions thereof can be used to map the location of the 50365 genes on a chromosome. This process is called chromosome mapping. Chromosome mapping is useful in correlating the 50365 sequences with genes associated with disease.

[247] Briefly, 50365 genes can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp in length) from the 50365 nucleotide sequences. These primers can then be used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the 50365 sequences will yield an amplified fragment.

[248] A panel of somatic cell hybrids in which each cell line contains either a single human chromosome or a small number of human chromosomes, and a full set of mouse chromosomes, can allow easy mapping of individual genes to specific human chromosomes (D'Eustachio P. *et al.* (1983) *Science* 220:919-924).

[249] Other mapping strategies e.g., *in situ* hybridization (described in Fan, Y. *et al.* (1990) *Proc. Natl. Acad. Sci. USA*, 87:6223-27), pre-screening with labeled flow-sorted chromosomes, and pre-selection by hybridization to chromosome specific cDNA libraries can be used to map 50365 to a chromosomal location.

[250] Fluorescence *in situ* hybridization (FISH) of a DNA sequence to a metaphase chromosomal spread can further be used to provide a precise chromosomal location in one step. The FISH technique can be used with a DNA sequence as short as 500 or 600 bases. However, clones larger than 1,000 bases have a higher likelihood of binding to a unique chromosomal location with sufficient signal intensity for simple detection. Preferably 1,000 bases, and more preferably 2,000 bases will suffice to get good results at a reasonable

amount of time. For a review of this technique, see Verma *et al.*, Human Chromosomes: A Manual of Basic Techniques ((1988) Pergamon Press, New York).

[251] Reagents for chromosome mapping can be used individually to mark a single chromosome or a single site on that chromosome, or panels of reagents can be used for
5 marking multiple sites and/or multiple chromosomes. Reagents corresponding to noncoding regions of the genes actually are preferred for mapping purposes. Coding sequences are more likely to be conserved within gene families, thus increasing the chance of cross hybridizations during chromosomal mapping.

[252] Once a sequence has been mapped to a precise chromosomal location, the
10 physical position of the sequence on the chromosome can be correlated with genetic map data. (Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man, available on-line through Johns Hopkins University Welch Medical Library). The relationship between a gene and a disease, mapped to the same chromosomal region, can then be identified through linkage analysis (co-inheritance of physically adjacent genes),
15 described in, for example, Egeland, J. *et al.* (1987) *Nature*, 325:783-787.

[253] Moreover, differences in the DNA sequences between individuals affected and unaffected with a disease associated with the 50365 gene, can be determined. If a mutation is observed in some or all of the affected individuals but not in any unaffected individuals, then the mutation is likely to be the causative agent of the particular disease.
20 Comparison of affected and unaffected individuals generally involves first looking for structural alterations in the chromosomes, such as deletions or translocations that are visible from chromosome spreads or detectable using PCR based on that DNA sequence. Ultimately, complete sequencing of genes from several individuals can be performed to confirm the presence of a mutation and to distinguish mutations from polymorphisms.

25 Tissue Typing

[254] 50365 sequences can be used to identify individuals from biological samples using, e.g., restriction fragment length polymorphism (RFLP). In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, the fragments separated, e.g., in a Southern blot, and probed to yield bands for identification. The
30 sequences of the present invention are useful as additional DNA markers for RFLP (described in U.S. Patent 5,272,057).

[255] Furthermore, the sequences of the present invention can also be used to determine the actual base-by-base DNA sequence of selected portions of an individual's genome. Thus, the 50365 nucleotide sequences described herein can be used to prepare two PCR primers from the 5' and 3' ends of the sequences. These primers can then be used to
5 amplify an individual's DNA and subsequently sequence it. Panels of corresponding DNA sequences from individuals, prepared in this manner, can provide unique individual identifications, as each individual will have a unique set of such DNA sequences due to allelic differences.

[256] Allelic variation occurs to some degree in the coding regions of these
10 sequences, and to a greater degree in the noncoding regions. Each of the sequences described herein can, to some degree, be used as a standard against which DNA from an individual can be compared for identification purposes. Because greater numbers of polymorphisms occur in the noncoding regions, fewer sequences are necessary to differentiate individuals. The noncoding sequences of SEQ ID NO:1 can provide positive
15 individual identification with a panel of perhaps 10 to 1,000 primers which each yield a noncoding amplified sequence of 100 bases. If predicted coding sequences, such as those in SEQ ID NO:3 are used, a more appropriate number of primers for positive individual identification would be 500-2,000.

[257] If a panel of reagents from 50365 nucleotide sequences described herein is
20 used to generate a unique identification database for an individual, those same reagents can later be used to identify tissue from that individual. Using the unique identification database, positive identification of the individual, living or dead, can be made from extremely small tissue samples.

Use of Partial 50365 Sequences in Forensic Biology

25 [258] DNA-based identification techniques can also be used in forensic biology. To make such an identification, PCR technology can be used to amplify DNA sequences taken from very small biological samples such as tissues, e.g., hair or skin, or body fluids, e.g., blood, saliva, or semen found at a crime scene. The amplified sequence can then be compared to a standard, thereby allowing identification of the origin of the biological
30 sample.

[259] The sequences of the present invention can be used to provide polynucleotide reagents, e.g., PCR primers, targeted to specific loci in the human genome, which can

enhance the reliability of DNA-based forensic identifications by, for example, providing another "identification marker" (i.e. another DNA sequence that is unique to a particular individual). As mentioned above, actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments. Sequences targeted to noncoding regions of SEQ ID NO:1 (e.g., fragments derived from the noncoding regions of SEQ ID NO:1 having a length of at least 20 bases, preferably at least 30 bases) are particularly appropriate for this use.

[260] The 50365 nucleotide sequences described herein can further be used to provide polynucleotide reagents, e.g., labeled or labelable probes which can be used in, for example, an *in situ* hybridization technique, to identify a specific tissue. This can be very useful in cases where a forensic pathologist is presented with a tissue of unknown origin. Panels of such 50365 probes can be used to identify tissue by species and/or by organ type.

[261] In a similar fashion, these reagents, e.g., 50365 primers or probes can be used to screen tissue culture for contamination (i.e. screen for the presence of a mixture of different types of cells in a culture).

Predictive Medicine

[262] The present invention also pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, and monitoring clinical trials are used for prognostic (predictive) purposes to thereby treat an individual.

20 [263] Generally, the invention provides, a method of determining if a subject is at risk for a disorder related to a lesion in or the misexpression of a gene which encodes 50365.

[264] Such disorders include, e.g., a disorder associated with the misexpression of 50365; a disorder of the glycolytic control system.

[265] The method includes one or more of the following:

25 [266] detecting, in a tissue of the subject, the presence or absence of a mutation which affects the expression of the 50365 gene, or detecting the presence or absence of a mutation in a region which controls the expression of the gene, e.g., a mutation in the 5' control region;

[267] detecting, in a tissue of the subject, the presence or absence of a mutation which alters the structure of the 50365 gene;

30 [268] detecting, in a tissue of the subject, the misexpression of the 50365 gene, at the mRNA level, e.g., detecting a non-wild type level of a mRNA ;

[269] detecting, in a tissue of the subject, the misexpression of the gene, at the protein level, e.g., detecting a non-wild type level of a 50365 polypeptide.

[270] In preferred embodiments the method includes: ascertaining the existence of at least one of: a deletion of one or more nucleotides from the 50365 gene; an insertion of one or more nucleotides into the gene, a point mutation, e.g., a substitution of one or more nucleotides of the gene, a gross chromosomal rearrangement of the gene, e.g., a translocation, inversion, or deletion.

[271] For example, detecting the genetic lesion can include: (i) providing a probe/primer including an oligonucleotide containing a region of nucleotide sequence which hybridizes to a sense or antisense sequence from SEQ ID NO:1, or naturally occurring mutants thereof or 5' or 3' flanking sequences naturally associated with the 50365 gene; (ii) exposing the probe/primer to nucleic acid of the tissue; and detecting, by hybridization, e.g., *in situ* hybridization, of the probe/primer to the nucleic acid, the presence or absence of the genetic lesion.

[272] In preferred embodiments detecting the misexpression includes ascertaining the existence of at least one of: an alteration in the level of a messenger RNA transcript of the 50365 gene; the presence of a non-wild type splicing pattern of a messenger RNA transcript of the gene; or a non-wild type level of 50365.

[273] Methods of the invention can be used prenatally or to determine if a subject's offspring will be at risk for a disorder.

[274] In preferred embodiments the method includes determining the structure of a 50365 gene, an abnormal structure being indicative of risk for the disorder.

[275] In preferred embodiments the method includes contacting a sample from the subject with an antibody to the 50365 protein or a nucleic acid, which hybridizes specifically with the gene. These and other embodiments are discussed below.

Diagnostic and Prognostic Assays

[276] Diagnostic and prognostic assays of the invention include method for assessing the expression level of 50365 molecules and for identifying variations and mutations in the sequence of 50365 molecules.

[277] **Expression Monitoring and Profiling.** The presence, level, or absence of 50365 protein or nucleic acid in a biological sample can be evaluated by obtaining a biological sample from a test subject and contacting the biological sample with a compound.

or an agent capable of detecting 50365 protein or nucleic acid (e.g., mRNA, genomic DNA) that encodes 50365 protein such that the presence of 50365 protein or nucleic acid is detected in the biological sample. The term "biological sample" includes tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. A preferred biological sample is serum. The level of expression of the 50365 gene can be measured in a number of ways, including, but not limited to: measuring the mRNA encoded by the 50365 genes; measuring the amount of protein encoded by the 50365 genes; or measuring the activity of the protein encoded by the 50365 genes.

[278] The level of mRNA corresponding to the 50365 gene in a cell can be determined both by *in situ* and by *in vitro* formats.

[279] The isolated mRNA can be used in hybridization or amplification assays that include, but are not limited to, Southern or Northern analyses, polymerase chain reaction analyses and probe arrays. One preferred diagnostic method for the detection of mRNA levels involves contacting the isolated mRNA with a nucleic acid molecule (probe) that can hybridize to the mRNA encoded by the gene being detected. The nucleic acid probe can be, for example, a full-length 50365 nucleic acid, such as the nucleic acid of SEQ ID NO:1, or a portion thereof, such as an oligonucleotide of at least 7, 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to 50365 mRNA or genomic DNA. The probe can be disposed on an address of an array, e.g., an array described below. Other suitable probes for use in the diagnostic assays are described herein.

[280] In one format, mRNA (or cDNA) is immobilized on a surface and contacted with the probes, for example by running the isolated mRNA on an agarose gel and transferring the mRNA from the gel to a membrane, such as nitrocellulose. In an alternative format, the probes are immobilized on a surface and the mRNA (or cDNA) is contacted with the probes, for example, in a two-dimensional gene chip array described below. A skilled artisan can adapt known mRNA detection methods for use in detecting the level of mRNA encoded by the 50365 genes.

[281] The level of mRNA in a sample that is encoded by one of 50365 can be evaluated with nucleic acid amplification, e.g., by rtPCR (Mullis (1987) U.S. Patent No. 4,683,202), ligase chain reaction (Barany (1991) *Proc. Natl. Acad. Sci. USA* 88:189-193), self sustained sequence replication (Guatelli *et al.*, (1990) *Proc. Natl. Acad. Sci. USA* 87:1874-1878), transcriptional amplification system (Kwoh *et al.*, (1989), *Proc. Natl. Acad.*

Sci. USA 86:1173-1177), Q-Beta Replicase (Lizardi *et al.*, (1988) *Bio/Technology* 6:1197), rolling circle replication (Lizardi *et al.*, U.S. Patent No. 5,854,033) or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques known in the art. As used herein, amplification primers are defined as being a pair of

5 nucleic acid molecules that can anneal to 5' or 3' regions of a gene (plus and minus strands, respectively, or vice-versa) and contain a short region in between. In general, amplification primers are from about 10 to 30 nucleotides in length and flank a region from about 50 to 200 nucleotides in length. Under appropriate conditions and with appropriate reagents, such primers permit the amplification of a nucleic acid molecule comprising the nucleotide

10 sequence flanked by the primers.

[282] For *in situ* methods, a cell or tissue sample can be prepared/processed and immobilized on a support, typically a glass slide, and then contacted with a probe that can hybridize to mRNA that encodes the 50365 gene being analyzed.

[283] In another embodiment, the methods further contacting a control sample with

15 a compound or agent capable of detecting 50365 mRNA, or genomic DNA, and comparing the presence of 50365 mRNA or genomic DNA in the control sample with the presence of 50365 mRNA or genomic DNA in the test sample. In still another embodiment, serial analysis of gene expression, as described in U.S. Patent No. 5,695,937, is used to detect 50365 transcript levels.

[284] A variety of methods can be used to determine the level of protein encoded by 50365. In general, these methods include contacting an agent that selectively binds to the protein, such as an antibody with a sample, to evaluate the level of protein in the sample. In a preferred embodiment, the antibody bears a detectable label. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (e.g.,

25 Fab or F(ab')₂) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (i.e., physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with a detectable substance. Examples of detectable substances are provided herein.

[285] The detection methods can be used to detect 50365 protein in a biological sample *in vitro* as well as *in vivo*. *In vitro* techniques for detection of 50365 protein include enzyme linked immunosorbent assays (ELISAs), immunoprecipitations, immunofluorescence, enzyme immunoassay (EIA), radioimmunoassay (RIA), and Western

blot analysis. *In vivo* techniques for detection of 50365 protein include introducing into a subject a labeled anti-50365 antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques. In another embodiment, the sample is labeled, e.g., biotinylated and
5 then contacted to the antibody, e.g., an anti-50365 antibody positioned on an antibody array (as described below). The sample can be detected, e.g., with avidin coupled to a fluorescent label.

[286] In another embodiment, the methods further include contacting the control sample with a compound or agent capable of detecting 50365 protein, and comparing the
10 presence of 50365 protein in the control sample with the presence of 50365 protein in the test sample.

[287] The invention also includes kits for detecting the presence of 50365 in a biological sample. For example, the kit can include a compound or agent capable of detecting 50365 protein or mRNA in a biological sample; and a standard. The compound or
15 agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect 50365 protein or nucleic acid.

[288] For antibody-based kits, the kit can include: (1) a first antibody (e.g., attached to a solid support) which binds to a polypeptide corresponding to a marker of the invention; and, optionally, (2) a second, different antibody which binds to either the
20 polypeptide or the first antibody and is conjugated to a detectable agent.

[289] For oligonucleotide-based kits, the kit can include: (1) an oligonucleotide, e.g., a detectably labeled oligonucleotide, which hybridizes to a nucleic acid sequence encoding a polypeptide corresponding to a marker of the invention or (2) a pair of primers useful for amplifying a nucleic acid molecule corresponding to a marker of the invention.
25 The kit can also include a buffering agent, a preservative, or a protein stabilizing agent. The kit can also include components necessary for detecting the detectable agent (e.g., an enzyme or a substrate). The kit can also contain a control sample or a series of control samples which can be assayed and compared to the test sample contained. Each component
30 of the kit can be enclosed within an individual container and all of the various containers can be within a single package, along with instructions for interpreting the results of the assays performed using the kit.

[290] The diagnostic methods described herein can identify subjects having, or at risk of developing, a disease or disorder associated with misexpressed or aberrant or

unwanted 50365 expression or activity. As used herein, the term "unwanted" includes an unwanted phenomenon involved in a biological response such as faulty glycolytic activity, including faulty phosphorylation of aldo- and ketohexoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine) or deregulated cell proliferation.

5 [291] In one embodiment, a disease or disorder associated with aberrant or unwanted 50365 expression or activity is identified. A test sample is obtained from a subject and 50365 protein or nucleic acid (e.g., mRNA or genomic DNA) is evaluated, wherein the level, e.g., the presence or absence, of 50365 protein or nucleic acid is diagnostic for a subject having or at risk of developing a disease or disorder associated with
10 aberrant or unwanted 50365 expression or activity. As used herein, a "test sample" refers to a biological sample obtained from a subject of interest, including a biological fluid (e.g., serum), cell sample, or tissue.

[292] The prognostic assays described herein can be used to determine whether a subject can be administered an agent (e.g., an agonist, antagonist, peptidomimetic, protein,
15 peptide, nucleic acid, small molecule, or other drug candidate) to treat a disease or disorder associated with aberrant or unwanted 50365 expression or activity. For example, such methods can be used to determine whether a subject can be effectively treated with an agent for a cell having faulty glycolytic activity, including faulty phosphorylation of aldo- and ketohexoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine).

20 [293] In another aspect, the invention features a computer medium having a plurality of digitally encoded data records. Each data record includes a value representing the level of expression of 50365 in a sample, and a descriptor of the sample. The descriptor of the sample can be an identifier of the sample, a subject from which the sample was derived (e.g., a patient), a diagnosis, or a treatment (e.g., a preferred treatment). In a
25 preferred embodiment, the data record further includes values representing the level of expression of genes other than 50365 (e.g., other genes associated with a 50365-disorder, or other genes on an array). The data record can be structured as a table, e.g., a table that is part of a database such as a relational database (e.g., a SQL database of the Oracle or Sybase database environments).

30 [294] Also featured is a method of evaluating a sample. The method includes providing a sample, e.g., from the subject, and determining a gene expression profile of the sample, wherein the profile includes a value representing the level of 50365 expression. The method can further include comparing the value or the profile (i.e., multiple values) to a

- reference value or reference profile. The gene expression profile of the sample can be obtained by any of the methods described herein (e.g., by providing a nucleic acid from the sample and contacting the nucleic acid to an array). The method can be used to diagnose faulty glycolytic activity disorders, including faulty phosphorylation of aldo- and
- 5 ketohehexoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine) in a subject wherein an increase in 50365 expression is an indication that the subject has or is disposed to having faulty glycolytic activity, including faulty phosphorylation of aldo- and ketohehexoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine). The method can be used to monitor a treatment for faulty glycolytic activity, including faulty
- 10 phosphorylation of aldo- and ketohehexoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine) in a subject. For example, the gene expression profile can be determined for a sample from a subject undergoing treatment. The profile can be compared to a reference profile or to a profile obtained from the subject prior to treatment or prior to onset of the disorder (see, e.g., Golub *et al.* (1999) *Science* 286:531).
- 15 [295] In yet another aspect, the invention features a method of evaluating a test compound (see also, "Screening Assays", above). The method includes providing a cell and a test compound; contacting the test compound to the cell; obtaining a subject expression profile for the contacted cell; and comparing the subject expression profile to one or more reference profiles. The profiles include a value representing the level of 50365 expression.
- 20 In a preferred embodiment, the subject expression profile is compared to a target profile, e.g., a profile for a normal cell or for desired condition of a cell. The test compound is evaluated favorably if the subject expression profile is more similar to the target profile than an expression profile obtained from an uncontacted cell.
- [296] In another aspect, the invention features, a method of evaluating a subject.
- 25 The method includes: a) obtaining a sample from a subject, e.g., from a caregiver, e.g., a caregiver who obtains the sample from the subject; b) determining a subject expression profile for the sample. Optionally, the method further includes either or both of steps: c) comparing the subject expression profile to one or more reference expression profiles; and d) selecting the reference profile most similar to the subject reference profile. The subject
- 30 expression profile and the reference profiles include a value representing the level of 50365 expression. A variety of routine statistical measures can be used to compare two reference profiles. One possible metric is the length of the distance vector that is the difference

between the two profiles. Each of the subject and reference profile is represented as a multi-dimensional vector, wherein each dimension is a value in the profile.

[297] The method can further include transmitting a result to a caregiver. The result can be the subject expression profile, a result of a comparison of the subject
5 expression profile with another profile, a most similar reference profile, or a descriptor of any of the aforementioned. The result can be transmitted across a computer network, e.g., the result can be in the form of a computer transmission, e.g., a computer data signal embedded in a carrier wave.

[298] Also featured is a computer medium having executable code for effecting the
10 following steps: receive a subject expression profile; access a database of reference expression profiles; and either i) select a matching reference profile most similar to the subject expression profile or ii) determine at least one comparison score for the similarity of the subject expression profile to at least one reference profile. The subject expression profile, and the reference expression profiles each include a value representing the level of
15 50365 expression.

Arrays and Uses Thereof

[299] In another aspect, the invention features an array that includes a substrate having a plurality of addresses. At least one address of the plurality includes a capture probe that binds specifically to a 50365 molecule (e.g., a 50365 nucleic acid or a 50365
20 polypeptide). The array can have a density of at least than 10, 50, 100, 200, 500, 1,000, 2,000, or 10,000 or more addresses/cm², and ranges between. In a preferred embodiment, the plurality of addresses includes at least 10, 100, 500, 1,000, 5,000, 10,000, 50,000 addresses. In a preferred embodiment, the plurality of addresses includes equal to or less than 10, 100, 500, 1,000, 5,000, 10,000, or 50,000 addresses. The substrate can be a two-
25 dimensional substrate such as a glass slide, a wafer (e.g., silica or plastic), a mass spectroscopy plate, or a three-dimensional substrate such as a gel pad. Addresses in addition to address of the plurality can be disposed on the array.

[300] In a preferred embodiment, at least one address of the plurality includes a nucleic acid capture probe that hybridizes specifically to a 50365 nucleic acid, e.g., the sense
30 or anti-sense strand. In one preferred embodiment, a subset of addresses of the plurality of addresses has a nucleic acid capture probe for 50365. Each address of the subset can include a capture probe that hybridizes to a different region of a 50365 nucleic acid. In another

preferred embodiment, addresses of the subset include a capture probe for a 50365 nucleic acid. Each address of the subset is unique, overlapping, and complementary to a different variant of 50365 (e.g., an allelic variant, or all possible hypothetical variants). The array can be used to sequence 50365 by hybridization (see, e.g., U.S. Patent No. 5,695,940).

5 [301] An array can be generated by various methods, e.g., by photolithographic methods (see, e.g., U.S. Patent Nos. 5,143,854; 5,510,270; and 5,527,681), mechanical methods (e.g., directed-flow methods as described in U.S. Patent No. 5,384,261), pin-based methods (e.g., as described in U.S. Pat. No. 5,288,514), and bead-based techniques (e.g., as described in PCT US/93/04145).

10 [302] In another preferred embodiment, at least one address of the plurality includes a polypeptide capture probe that binds specifically to a 50365 polypeptide or fragment thereof. The polypeptide can be a naturally-occurring interaction partner of 50365 polypeptide. Preferably, the polypeptide is an antibody, e.g., an antibody described herein (see "Anti-50365 Antibodies," above), such as a monoclonal antibody or a single-chain
15 antibody.

[303] In another aspect, the invention features a method of analyzing the expression of 50365. The method includes providing an array as described above; contacting the array with a sample and detecting binding of a 50365-molecule (e.g., nucleic acid or polypeptide) to the array. In a preferred embodiment, the array is a nucleic acid array. Optionally the
20 method further includes amplifying nucleic acid from the sample prior or during contact with the array.

[304] In another embodiment, the array can be used to assay gene expression in a tissue to ascertain tissue specificity of genes in the array, particularly the expression of 50365. If a sufficient number of diverse samples is analyzed, clustering (e.g.,
25 hierarchical clustering, k-means clustering, Bayesian clustering and the like) can be used to identify other genes which are co-regulated with 50365. For example, the array can be used for the quantitation of the expression of multiple genes. Thus, not only tissue specificity, but also the level of expression of a battery of genes in the tissue is ascertained. Quantitative data can be used to group (e.g., cluster) genes on the basis of their tissue
30 expression *per se* and level of expression in that tissue.

[305] For example, array analysis of gene expression can be used to assess the effect of cell-cell interactions on 50365 expression. A first tissue can be perturbed and nucleic acid from a second tissue that interacts with the first tissue can be analyzed. In this

context, the effect of one cell type on another cell type in response to a biological stimulus can be determined, e.g., to monitor the effect of cell-cell interaction at the level of gene expression.

[306] In another embodiment, cells are contacted with a therapeutic agent. The expression profile of the cells is determined using the array, and the expression profile is compared to the profile of like cells not contacted with the agent. For example, the assay can be used to determine or analyze the molecular basis of an undesirable effect of the therapeutic agent. If an agent is administered therapeutically to treat one cell type but has an undesirable effect on another cell type, the invention provides an assay to determine the molecular basis of the undesirable effect and thus provides the opportunity to co-administer a counteracting agent or otherwise treat the undesired effect. Similarly, even within a single cell type, undesirable biological effects can be determined at the molecular level. Thus, the effects of an agent on expression of other than the target gene can be ascertained and counteracted.

[307] In another embodiment, the array can be used to monitor expression of one or more genes in the array with respect to time. For example, samples obtained from different time points can be probed with the array. Such analysis can identify and/or characterize the development of a 50365-associated disease or disorder, and processes, such as a cellular transformation associated with a 50365-associated disease or disorder. The method can also evaluate the treatment and/or progression of a 50365-associated disease or disorder

[308] The array is also useful for ascertaining differential expression patterns of one or more genes in normal and abnormal cells. This provides a battery of genes (e.g., including 50365) that could serve as a molecular target for diagnosis or therapeutic intervention.

[309] In another aspect, the invention features an array having a plurality of addresses. Each address of the plurality includes a unique polypeptide. At least one address of the plurality has disposed thereon a 50365 polypeptide or fragment thereof. Methods of producing polypeptide arrays are described in the art, e.g., in De Wildt *et al.* (2000). *Nature Biotech.* 18, 989-994; Lueking *et al.* (1999). *Anal. Biochem.* 270, 103-111; Ge, H. (2000). *Nucleic Acids Res.* 28, e3, I-VII; MacBeath, G., and Schreiber, S.L. (2000). *Science* 289, 1760-1763; and WO 99/51773A1. In a preferred embodiment, each addresses of the plurality has disposed thereon a polypeptide at least 60, 70, 80, 85, 90, 95 or 99 % identical to a 50365 polypeptide or fragment thereof. For example, multiple variants of a 50365

polypeptide (e.g., encoded by allelic variants, site-directed mutants, random mutants, or combinatorial mutants) can be disposed at individual addresses of the plurality. Addresses in addition to the address of the plurality can be disposed on the array.

[310] The polypeptide array can be used to detect a 50365 binding compound, e.g.,
5 an antibody in a sample from a subject with specificity for a 50365 polypeptide or the presence of a 50365-binding protein or ligand.

[311] The array is also useful for ascertaining the effect of the expression of a gene on the expression of other genes in the same cell or in different cells (e.g., ascertaining the effect of 50365 expression on the expression of other genes). This provides, for example,
10 for a selection of alternate molecular targets for therapeutic intervention if the ultimate or downstream target cannot be regulated.

[312] In another aspect, the invention features a method of analyzing a plurality of probes. The method is useful, e.g., for analyzing gene expression. The method includes: providing a two dimensional array having a plurality of addresses, each address of the
15 plurality being positionally distinguishable from each other address of the plurality having a unique capture probe, e.g., wherein the capture probes are from a cell or subject which express 50365 or from a cell or subject in which a 50365 mediated response has been elicited, e.g., by contact of the cell with 50365 nucleic acid or protein, or administration to the cell or subject 50365 nucleic acid or protein; providing a two dimensional array having a
20 plurality of addresses, each address of the plurality being positionally distinguishable from each other address of the plurality, and each address of the plurality having a unique capture probe, e.g., wherein the capture probes are from a cell or subject which does not express 50365 (or does not express as highly as in the case of the 50365 positive plurality of capture probes) or from a cell or subject in which a 50365 mediated response has not been
25 elicited (or has been elicited to a lesser extent than in the first sample); contacting the array with one or more inquiry probes (which is preferably other than a 50365 nucleic acid, polypeptide, or antibody), and thereby evaluating the plurality of capture probes. Binding, e.g., in the case of a nucleic acid, hybridization with a capture probe at an address of the plurality, is detected, e.g., by signal generated from a label attached to the nucleic acid,
30 polypeptide, or antibody.

[313] In another aspect, the invention features a method of analyzing a plurality of probes or a sample. The method is useful, e.g., for analyzing gene expression. The method includes: providing a two dimensional array having a plurality of addresses, each address of

the plurality being positionally distinguishable from each other address of the plurality having a unique capture probe, contacting the array with a first sample from a cell or subject which express or mis-express 50365 or from a cell or subject in which a 50365-mediated response has been elicited, e.g., by contact of the cell with 50365 nucleic acid or protein, or administration to the cell or subject 50365 nucleic acid or protein; providing a two dimensional array having a plurality of addresses, each address of the plurality being positionally distinguishable from each other address of the plurality, and each address of the plurality having a unique capture probe, and contacting the array with a second sample from a cell or subject which does not express 50365 (or does not express as highly as in the case of the 50365 positive plurality of capture probes) or from a cell or subject which in which a 50365 mediated response has not been elicited (or has been elicited to a lesser extent than in the first sample); and comparing the binding of the first sample with the binding of the second sample. Binding, e.g., in the case of a nucleic acid, hybridization with a capture probe at an address of the plurality, is detected, e.g., by signal generated from a label attached to the nucleic acid, polypeptide, or antibody. The same array can be used for both samples or different arrays can be used. If different arrays are used the plurality of addresses with capture probes should be present on both arrays.

[314] In another aspect, the invention features a method of analyzing 50365, e.g., analyzing structure, function, or relatedness to other nucleic acid or amino acid sequences. The method includes: providing a 50365 nucleic acid or amino acid sequence; comparing the 50365 sequence with one or more preferably a plurality of sequences from a collection of sequences, e.g., a nucleic acid or protein sequence database; to thereby analyze 50365.

Detection of Variations or Mutations

[315] The methods of the invention can also be used to detect genetic alterations in a 50365 gene, thereby determining if a subject with the altered gene is at risk for a disorder characterized by misregulation in 50365 protein activity or nucleic acid expression, such as faulty glycolytic activity, including faulty phosphorylation of aldo- and ketohexoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine). In preferred embodiments, the methods include detecting, in a sample from the subject, the presence or absence of a genetic alteration characterized by at least one of an alteration affecting the integrity of a gene encoding a 50365-protein, or the mis-expression of the 50365 gene. For example, such genetic alterations can be detected by ascertaining the existence of at least one of 1) a

deletion of one or more nucleotides from a 50365 gene; 2) an addition of one or more nucleotides to a 50365 gene; 3) a substitution of one or more nucleotides of a 50365 gene, 4) a chromosomal rearrangement of a 50365 gene; 5) an alteration in the level of a messenger RNA transcript of a 50365 gene, 6) aberrant modification of a 50365 gene, such as of the methylation pattern of the genomic DNA, 7) the presence of a non-wild type splicing pattern of a messenger RNA transcript of a 50365 gene, 8) a non-wild type level of a 50365-protein, 9) allelic loss of a 50365 gene, and 10) inappropriate post-translational modification of a 50365-protein.

[316] An alteration can be detected without a probe/primer in a polymerase chain reaction, such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR), the latter of which can be particularly useful for detecting point mutations in the 50365-gene. This method can include the steps of collecting a sample of cells from a subject, isolating nucleic acid (e.g., genomic, mRNA or both) from the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a 50365 gene under conditions such that hybridization and amplification of the 50365-gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. It is anticipated that PCR and/or LCR may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein. Alternatively, other amplification methods described herein or known in the art can be used.

[317] In another embodiment, mutations in a 50365 gene from a sample cell can be identified by detecting alterations in restriction enzyme cleavage patterns. For example, sample and control DNA is isolated, amplified (optionally), digested with one or more restriction endonucleases, and fragment length sizes are determined, e.g., by gel electrophoresis and compared. Differences in fragment length sizes between sample and control DNA indicates mutations in the sample DNA. Moreover, the use of sequence specific ribozymes (see, for example, U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

[318] In other embodiments, genetic mutations in 50365 can be identified by hybridizing a sample and control nucleic acids, e.g., DNA or RNA, two-dimensional arrays, e.g., chip based arrays. Such arrays include a plurality of addresses, each of which is positionally distinguishable from the other. A different probe is located at each address of the plurality. A probe can be complementary to a region of a 50365 nucleic acid or a

- putative variant (e.g., allelic variant) thereof. A probe can have one or more mismatches to a region of a 50365 nucleic acid (e.g., a destabilizing mismatch). The arrays can have a high density of addresses, e.g., can contain hundreds or thousands of oligonucleotides probes (Cronin, M.T. *et al.* (1996) *Human Mutation* 7: 244-255; Kozal, M.J. *et al.* (1996) *Nature Medicine* 2: 753-759). For example, genetic mutations in 50365 can be identified in two-dimensional arrays containing light-generated DNA probes as described in Cronin, M.T. *et al. supra*. Briefly, a first hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base changes between the sequences by making linear arrays of sequential overlapping probes. This step allows the identification of point mutations. This step is followed by a second hybridization array that allows the characterization of specific mutations by using smaller, specialized probe arrays complementary to all variants or mutations detected. Each mutation array is composed of parallel probe sets, one complementary to the wild-type gene and the other complementary to the mutant gene.
- 15 [319] In yet another embodiment, any of a variety of sequencing reactions known in the art can be used to directly sequence the 50365 gene and detect mutations by comparing the sequence of the sample 50365 with the corresponding wild-type (control) sequence. Automated sequencing procedures can be utilized when performing the diagnostic assays ((1995) *Biotechniques* 19:448), including sequencing by mass spectrometry.
- 20 [320] Other methods for detecting mutations in the 50365 gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA heteroduplexes (Myers *et al.* (1985) *Science* 230:1242; Cotton *et al.* (1988) *Proc. Natl Acad Sci USA* 85:4397; Saleeba *et al.* (1992) *Methods Enzymol.* 217:286-295).
- 25 [321] In still another embodiment, the mismatch cleavage reaction employs one or more proteins that recognize mismatched base pairs in double-stranded DNA (so called "DNA mismatch repair" enzymes) in defined systems for detecting and mapping point mutations in 50365 cDNAs obtained from samples of cells. For example, the mutY enzyme of *E. coli* cleaves A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches (Hsu *et al.* (1994) *Carcinogenesis* 15:1657-1662; U.S. Patent No. 5,459,039).
- 30 [322] In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in 50365 genes. For example, single strand conformation polymorphism

(SSCP) may be used to detect differences in electrophoretic mobility between mutant and wild type nucleic acids (Orita *et al.* (1989) *Proc Natl. Acad. Sci USA*: 86:2766, see also Cotton (1993) *Mutat. Res.* 285:125-144; and Hayashi (1992) *Genet. Anal. Tech. Appl* 9:73-79). Single-stranded DNA fragments of sample and control 50365 nucleic acids will be
5 denatured and allowed to renature. The secondary structure of single-stranded nucleic acids varies according to sequence, the resulting alteration in electrophoretic mobility enables the detection of even a single base change. The DNA fragments may be labeled or detected with labeled probes. The sensitivity of the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In a
10 preferred embodiment, the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex molecules on the basis of changes in electrophoretic mobility (Keen *et al.* (1991) *Trends Genet* 7:5).

[323] In yet another embodiment, the movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing
15 gradient gel electrophoresis (DGGE) (Myers *et al.* (1985) *Nature* 313:495). When DGGE is used as the method of analysis, DNA will be modified to insure that it does not completely denature, for example by adding a GC clamp of approximately 40 bp of high-melting GC-rich DNA by PCR. In a further embodiment, a temperature gradient is used in place of a denaturing gradient to identify differences in the mobility of control and sample DNA
20 (Rosenbaum and Reissner (1987) *Biophys Chem* 265:12753).

[324] Examples of other techniques for detecting point mutations include, but are not limited to, selective oligonucleotide hybridization, selective amplification, or selective primer extension (Saiki *et al.* (1986) *Nature* 324:163); Saiki *et al.* (1989) *Proc. Natl Acad. Sci USA* 86:6230). A further method of detecting point mutations is the chemical ligation of
25 oligonucleotides as described in Xu *et al.* ((2001) *Nature Biotechnol.* 19:148). Adjacent oligonucleotides, one of which selectively anneals to the query site, are ligated together if the nucleotide at the query site of the sample nucleic acid is complementary to the query oligonucleotide; ligation can be monitored, e.g., by fluorescent dyes coupled to the oligonucleotides.

30 [325] Alternatively, allele specific amplification technology that depends on selective PCR amplification may be used in conjunction with the instant invention. Oligonucleotides used as primers for specific amplification may carry the mutation of interest in the center of the molecule (so that amplification depends on differential

hybridization) (Gibbs *et al.* (1989) *Nucleic Acids Res.* 17:2437-2448) or at the extreme 3' end of one primer where, under appropriate conditions, mismatch can prevent, or reduce polymerase extension (Prossner (1993) *Tibtech* 11:238). In addition it may be desirable to introduce a novel restriction site in the region of the mutation to create cleavage-based detection (Gasparini *et al.* (1992) *Mol. Cell Probes* 6:1). It is anticipated that in certain 5 embodiments amplification may also be performed using Taq ligase for amplification (Barany (1991) *Proc. Natl. Acad. Sci USA* 88:189). In such cases, ligation will occur only if there is a perfect match at the 3' end of the 5' sequence making it possible to detect the presence of a known mutation at a specific site by looking for the presence or absence of 10 amplification.

[326] In another aspect, the invention features a set of oligonucleotides. The set includes a plurality of oligonucleotides, each of which is at least partially complementary (e.g., at least 50%, 60%, 70%, 80%, 90%, 92%, 95%, 97%, 98%, or 99% complementary) to a 50365 nucleic acid.

15 [327] In a preferred embodiment the set includes a first and a second oligonucleotide. The first and second oligonucleotide can hybridize to the same or to different locations of SEQ ID NO:1 or the complement of SEQ ID NO:1. Different locations can be different but overlapping, or non-overlapping on the same strand. The first and second oligonucleotide can hybridize to sites on the same or on different strands.

20 [328] The set can be useful, e.g., for identifying SNP's, or identifying specific alleles of 50365. In a preferred embodiment, each oligonucleotide of the set has a different nucleotide at an interrogation position. In one embodiment, the set includes two oligonucleotides, each complementary to a different allele at a locus, e.g., a biallelic or polymorphic locus.

25 [329] In another embodiment, the set includes four oligonucleotides, each having a different nucleotide (e.g., adenine, guanine, cytosine, or thymidine) at the interrogation position. The interrogation position can be a SNP or the site of a mutation. In another preferred embodiment, the oligonucleotides of the plurality are identical in sequence to one another (except for differences in length). The oligonucleotides can be provided with 30 differential labels, such that an oligonucleotide that hybridizes to one allele provides a signal that is distinguishable from an oligonucleotide that hybridizes to a second allele. In still another embodiment, at least one of the oligonucleotides of the set has a nucleotide change at a position in addition to a query position, e.g., a destabilizing mutation to decrease the T_m

of the oligonucleotide. In another embodiment, at least one oligonucleotide of the set has a non-natural nucleotide, e.g., inosine. In a preferred embodiment, the oligonucleotides are attached to a solid support, e.g., to different addresses of an array or to different beads or nanoparticles.

5 [330] In a preferred embodiment the set of oligo nucleotides can be used to specifically amplify, e.g., by PCR, or detect, a 50365 nucleic acid.

[331] The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits comprising at least one probe nucleic acid or antibody reagent described herein, which may be conveniently used, e.g., in clinical settings to diagnose
10 patients exhibiting symptoms or family history of a disease or illness involving a 50365 gene.

Use of 50365 Molecules as Surrogate Markers

[332] The 50365 molecules of the invention are also useful as markers of disorders or disease states, as markers for precursors of disease states, as markers for predisposition of
15 disease states, as markers of drug activity, or as markers of the pharmacogenomic profile of a subject. Using the methods described herein, the presence, absence and/or quantity of the 50365 molecules of the invention may be detected, and may be correlated with one or more biological states in vivo. For example, the 50365 molecules of the invention may serve as surrogate markers for one or more disorders or disease states or for conditions leading up to
20 disease states. As used herein, a "surrogate marker" is an objective biochemical marker which correlates with the absence or presence of a disease or disorder, or with the progression of a disease or disorder (e.g., with the presence or absence of a tumor). The presence or quantity of such markers is independent of the disease. Therefore, these markers may serve to indicate whether a particular course of treatment is effective in
25 lessening a disease state or disorder. Surrogate markers are of particular use when the presence or extent of a disease state or disorder is difficult to assess through standard methodologies (e.g., early stage tumors), or when an assessment of disease progression is desired before a potentially dangerous clinical endpoint is reached (e.g., an assessment of cardiovascular disease may be made using cholesterol levels as a surrogate marker, and an
30 analysis of HIV infection may be made using HIV RNA levels as a surrogate marker, well in advance of the undesirable clinical outcomes of myocardial infarction or fully-developed

AIDS). Examples of the use of surrogate markers in the art include: Koomen *et al.* (2000) *J. Mass. Spectrom.* 35: 258-264; and James (1994) *AIDS Treatment News Archive* 209.

[333] The 50365 molecules of the invention are also useful as pharmacodynamic markers. As used herein, a "pharmacodynamic marker" is an objective biochemical marker which correlates specifically with drug effects. The presence or quantity of a pharmacodynamic marker is not related to the disease state or disorder for which the drug is being administered; therefore, the presence or quantity of the marker is indicative of the presence or activity of the drug in a subject. For example, a pharmacodynamic marker may be indicative of the concentration of the drug in a biological tissue, in that the marker is either expressed or transcribed or not expressed or transcribed in that tissue in relationship to the level of the drug. In this fashion, the distribution or uptake of the drug may be monitored by the pharmacodynamic marker. Similarly, the presence or quantity of the pharmacodynamic marker may be related to the presence or quantity of the metabolic product of a drug, such that the presence or quantity of the marker is indicative of the relative breakdown rate of the drug in vivo. Pharmacodynamic markers are of particular use in increasing the sensitivity of detection of drug effects, particularly when the drug is administered in low doses. Since even a small amount of a drug may be sufficient to activate multiple rounds of marker (e.g., a 50365 marker) transcription or expression, the amplified marker may be in a quantity which is more readily detectable than the drug itself. Also, the marker may be more easily detected due to the nature of the marker itself; for example, using the methods described herein, anti-50365 antibodies may be employed in an immune-based detection system for a 50365 protein marker, or 50365-specific radiolabeled probes may be used to detect a 50365 mRNA marker. Furthermore, the use of a pharmacodynamic marker may offer mechanism-based prediction of risk due to drug treatment beyond the range of possible direct observations. Examples of the use of pharmacodynamic markers in the art include: Matsuda *et al.* US 6,033,862; Hattis *et al.* (1991) *Env. Health Perspect.* 90: 229-238; Schentag (1999) *Am. J. Health-Syst. Pharm.* 56 Suppl. 3: S21-S24; and Nicolau (1999) *Am. J. Health-Syst. Pharm.* 56 Suppl. 3: S16-S20.

[334] The 50365 molecules of the invention are also useful as pharmacogenomic markers. As used herein, a "pharmacogenomic marker" is an objective biochemical marker which correlates with a specific clinical drug response or susceptibility in a subject (see, e.g., McLeod *et al.* (1999) *Eur. J. Cancer* 35:1650-1652). The presence or quantity of the pharmacogenomic marker is related to the predicted response of the subject to a specific

drug or class of drugs prior to administration of the drug. By assessing the presence or quantity of one or more pharmacogenomic markers in a subject, a drug therapy which is most appropriate for the subject, or which is predicted to have a greater degree of success, may be selected. For example, based on the presence or quantity of RNA, or protein (e.g., 50365 protein or RNA) for specific tumor markers in a subject, a drug or course of treatment may be selected that is optimized for the treatment of the specific tumor likely to be present in the subject. Similarly, the presence or absence of a specific sequence mutation in 50365 DNA may correlate 50365 drug response. The use of pharmacogenomic markers therefore permits the application of the most appropriate treatment for each subject without having to administer the therapy.

Pharmaceutical Compositions

[335] The nucleic acid and polypeptides, fragments thereof, as well as anti-50365 antibodies (also referred to herein as "active compounds") of the invention can be incorporated into pharmaceutical compositions. Such compositions typically include the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein the language "pharmaceutically acceptable carrier" includes solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Supplementary active compounds can also be incorporated into the compositions.

[336] A pharmaceutical composition is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

- [337] Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It should be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.
- 20 [338] Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those
- 25 enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.
- [339] Oral compositions generally include an inert diluent or an edible carrier. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules, e.g., gelatin capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as
- 30

part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

[340] For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

[341] Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

[342] The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

[343] In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

[344] It is advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein

refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier.

[345] Toxicity and therapeutic efficacy of such compounds can be determined by
5 standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds which exhibit high therapeutic indices are preferred. While compounds that
10 exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

[346] The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies
15 preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma
20 concentration range that includes the IC50 (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[347] As defined herein, a therapeutically effective amount of protein or
25 polypeptide (*i.e.*, an effective dosage) ranges from about 0.001 to 30 mg/kg body weight, preferably about 0.01 to 25 mg/kg body weight, more preferably about 0.1 to 20 mg/kg body weight, and even more preferably about 1 to 10 mg/kg, 2 to 9 mg/kg, 3 to 8 mg/kg, 4 to 7 mg/kg, or 5 to 6 mg/kg body weight. The protein or polypeptide can be administered one time per week for between about 1 to 10 weeks, preferably between 2 to 8 weeks, more
30 preferably between about 3 to 7 weeks, and even more preferably for about 4, 5, or 6 weeks. The skilled artisan will appreciate that certain factors may influence the dosage and timing required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other

diseases present. Moreover, treatment of a subject with a therapeutically effective amount of a protein, polypeptide, or antibody can include a single treatment or, preferably, can include a series of treatments.

[348] For antibodies, the preferred dosage is 0.1 mg/kg of body weight (generally 5 10 mg/kg to 20 mg/kg). If the antibody is to act in the brain, a dosage of 50 mg/kg to 100 mg/kg is usually appropriate. Generally, partially human antibodies and fully human antibodies have a longer half-life within the human body than other antibodies. Accordingly, lower dosages and less frequent administration is often possible. Modifications such as lipidation can be used to stabilize antibodies and to enhance uptake 10 and tissue penetration (e.g., into the brain). A method for lipidation of antibodies is described by Cruikshank *et al.* ((1997) *J. Acquired Immune Deficiency Syndromes and Human Retrovirology* 14:193).

[349] The present invention encompasses agents which modulate expression or activity. An agent may, for example, be a small molecule. For example, such small 15 molecules include, but are not limited to, peptides, peptidomimetics (e.g., peptoids), amino acids, amino acid analogs, polynucleotides, polynucleotide analogs, nucleotides, nucleotide analogs, organic or inorganic compounds (i.e., including heteroorganic and organometallic compounds) having a molecular weight less than about 10,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 5,000 grams per mole, 20 organic or inorganic compounds having a molecular weight less than about 1,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 500 grams per mole, and salts, esters, and other pharmaceutically acceptable forms of such compounds.

[350] Exemplary doses include milligram or microgram amounts of the small molecule per kilogram of subject or sample weight (e.g., about 1 microgram per kilogram to 25 about 500 milligrams per kilogram, about 100 micrograms per kilogram to about 5 milligrams per kilogram, or about 1 microgram per kilogram to about 50 micrograms per kilogram. It is furthermore understood that appropriate doses of a small molecule depend upon the potency of the small molecule with respect to the expression or activity to be modulated. When one or more of these small molecules is to be administered to an animal 30 (e.g., a human) in order to modulate expression or activity of a polypeptide or nucleic acid of the invention, a physician, veterinarian, or researcher may, for example, prescribe a relatively low dose at first, subsequently increasing the dose until an appropriate response is obtained. In addition, it is understood that the specific dose level for any particular animal

subject will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, gender, and diet of the subject, the time of administration, the route of administration, the rate of excretion, any drug combination, and the degree of expression or activity to be modulated.

- 5 [351] An antibody (or fragment thereof) may be conjugated to a therapeutic moiety such as a cytotoxin, a therapeutic agent or a radioactive metal ion. A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, 10 mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine 15 (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine).
- 20 [352] The conjugates of the invention can be used for modifying a given biological response, the drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor, α - 25 interferon, β -interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator; or, biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors.
- 30 [353] Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described by Segal in U.S. Patent No. 4,676,980.
- [354] The nucleic acid molecules of the invention can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for

example, intravenous injection, local administration (see U.S. Patent 5,328,470) or by stereotactic injection (see e.g., Chen *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, e.g., retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system.

[355] The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

10 Methods of Treatment

[356] The present invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disorder or having a disorder associated with aberrant or unwanted 50365 expression or activity. As used herein, the term "treatment" is defined as the application or administration of a therapeutic agent to a patient, or application or administration of a therapeutic agent to an isolated tissue or cell line from a patient, who has a disease, a symptom of disease or a predisposition toward a disease, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve or affect the disease, the symptoms of disease or the predisposition toward disease. A therapeutic agent includes, but is not limited to, small molecules, peptides, antibodies, ribozymes and antisense oligonucleotides.

[357] With regards to both prophylactic and therapeutic methods of treatment, such treatments may be specifically tailored or modified, based on knowledge obtained from the field of pharmacogenomics. "Pharmacogenomics", as used herein, refers to the application of genomics technologies such as gene sequencing, statistical genetics, and gene expression analysis to drugs in clinical development and on the market. More specifically, the term refers the study of how a patient's genes determine his or her response to a drug (e.g., a patient's "drug response phenotype", or "drug response genotype".) Thus, another aspect of the invention provides methods for tailoring an individual's prophylactic or therapeutic treatment with either the 50365 molecules of the present invention or 50365 modulators according to that individual's drug response genotype. Pharmacogenomics allows a clinician or physician to target prophylactic or therapeutic treatments to patients who will most

benefit from the treatment and to avoid treatment of patients who will experience toxic drug-related side effects.

[358] In one aspect, the invention provides a method for preventing in a subject, a disease or condition associated with an aberrant or unwanted 50365 expression or activity, by administering to the subject a 50365 or an agent which modulates 50365 expression or at least one 50365 activity. Subjects at risk for a disease which is caused or contributed to by aberrant or unwanted 50365 expression or activity can be identified by, for example, any or a combination of diagnostic or prognostic assays as described herein. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the 50365 aberrance, such that a disease or disorder is prevented or, alternatively, delayed in its progression. Depending on the type of 50365 aberrance, for example, a 50365, 50365 agonist or 50365 antagonist agent can be used for treating the subject. The appropriate agent can be determined based on screening assays described herein.

[359] It is possible that some 50365 disorders can be caused, at least in part, by an abnormal level of gene product, or by the presence of a gene product exhibiting abnormal activity. As such, the reduction in the level and/or activity of such gene products would bring about the amelioration of disorder symptoms.

[360] The 50365 molecules can act as novel diagnostic targets and therapeutic agents for controlling one or more of cellular proliferative and/or differentiative disorders as described above, as well as disorders of the colon, pulmonary disorders, disorders associated with bone metabolism, immune disorders (e.g., inflammatory disorders), cardiovascular disorders, liver disorders, viral diseases, pain or metabolic disorders.

[361] Disorders involving the colon include, but are not limited to, congenital anomalies, such as atresia and stenosis, Meckel diverticulum, congenital aganglionic megacolon-Hirschsprung disease; enterocolitis, such as diarrhea and dysentery, infectious enterocolitis, including viral gastroenteritis, bacterial enterocolitis, necrotizing enterocolitis, antibiotic-associated colitis (pseudomembranous colitis), and collagenous and lymphocytic colitis, miscellaneous intestinal inflammatory disorders, including parasites and protozoa, acquired immunodeficiency syndrome, transplantation, drug-induced intestinal injury, radiation enterocolitis, neutropenic colitis (typhlitis), and diversion colitis; idiopathic inflammatory bowel disease, such as Crohn disease and ulcerative colitis; tumors of the colon, such as non-neoplastic polyps, adenomas, familial syndromes, colorectal carcinogenesis, colorectal carcinoma, and carcinoid tumors.

[362] Disorders involving the liver include, but are not limited to, hepatic injury; jaundice and cholestasis, such as bilirubin and bile formation; hepatic failure and cirrhosis, such as cirrhosis, portal hypertension, including ascites, portosystemic shunts, and splenomegaly; infectious disorders, such as viral hepatitis, including hepatitis A-E infection and infection by other hepatitis viruses, clinicopathologic syndromes, such as the carrier state, asymptomatic infection, acute viral hepatitis, chronic viral hepatitis, and fulminant hepatitis; autoimmune hepatitis; drug- and toxin-induced liver disease, such as alcoholic liver disease; inborn errors of metabolism and pediatric liver disease, such as hemochromatosis, Wilson disease, α_1 -antitrypsin deficiency, and neonatal hepatitis; intrahepatic biliary tract disease, such as secondary biliary cirrhosis, primary biliary cirrhosis, primary sclerosing cholangitis, and anomalies of the biliary tree; circulatory disorders, such as impaired blood flow into the liver, including hepatic artery compromise and portal vein obstruction and thrombosis, impaired blood flow through the liver, including passive congestion and centrilobular necrosis and peliosis hepatis, hepatic vein outflow obstruction, including hepatic vein thrombosis (Budd-Chiari syndrome) and veno-occlusive disease; hepatic disease associated with pregnancy, such as preeclampsia and eclampsia, acute fatty liver of pregnancy, and intrahepatic cholestasis of pregnancy; hepatic complications of organ or bone marrow transplantation, such as drug toxicity after bone marrow transplantation, graft-versus-host disease and liver rejection, and nonimmunologic damage to liver allografts; tumors and tumorous conditions, such as nodular hyperplasias, adenomas, and malignant tumors, including primary carcinoma of the liver and metastatic tumors.

[363] Examples of disorders of the lung include, but are not limited to, congenital anomalies; atelectasis; diseases of vascular origin, such as pulmonary congestion and edema, including hemodynamic pulmonary edema and edema caused by microvascular injury, adult respiratory distress syndrome (diffuse alveolar damage), pulmonary embolism, hemorrhage, and infarction, and pulmonary hypertension and vascular sclerosis; chronic obstructive pulmonary disease, such as emphysema, chronic bronchitis, bronchial asthma, and bronchiectasis; diffuse interstitial (infiltrative, restrictive) diseases, such as pneumoconioses, sarcoidosis, idiopathic pulmonary fibrosis, desquamative interstitial pneumonitis, hypersensitivity pneumonitis, pulmonary eosinophilia (pulmonary infiltration with eosinophilia), Bronchiolitis obliterans-organizing pneumonia, diffuse pulmonary hemorrhage syndromes, including Goodpasture syndrome, idiopathic pulmonary

hemosiderosis and other hemorrhagic syndromes, pulmonary involvement in collagen vascular disorders, and pulmonary alveolar proteinosis; complications of therapies, such as drug-induced lung disease, radiation-induced lung disease, and lung transplantation; tumors, such as bronchogenic carcinoma, including paraneoplastic syndromes, bronchioloalveolar carcinoma, neuroendocrine tumors, such as bronchial carcinoid, miscellaneous tumors, and metastatic tumors; pathologies of the pleura, including inflammatory pleural effusions, noninflammatory pleural effusions, pneumothorax, and pleural tumors, including solitary fibrous tumors (pleural fibroma) and malignant mesothelioma.

[364] Examples of proliferative disorders, in addition to those mentioned above, include hematopoietic neoplastic disorders. As used herein, the term "hematopoietic neoplastic disorders" includes diseases involving hyperplastic/neoplastic cells of hematopoietic origin, e.g., arising from myeloid, lymphoid or erythroid lineages, or precursor cells thereof. Preferably, the diseases arise from poorly differentiated acute leukemias, e.g., erythroblastic leukemia and acute megakaryoblastic leukemia. Additional exemplary myeloid disorders include, but are not limited to, acute promyeloid leukemia (APML), acute myelogenous leukemia (AML) and chronic myelogenous leukemia (CML) (reviewed in Vaickus, L. (1991) *Crit Rev. in Oncol./Hematol.* 11:267-97); lymphoid malignancies include, but are not limited to acute lymphoblastic leukemia (ALL) which includes B-lineage ALL and T-lineage ALL, chronic lymphocytic leukemia (CLL), prolymphocytic leukemia (PLL), hairy cell leukemia (HLL) and Waldenstrom's macroglobulinemia (WM). Additional forms of malignant lymphomas include, but are not limited to non-Hodgkin lymphoma and variants thereof, peripheral T cell lymphomas, adult T cell leukemia/lymphoma (ATL), cutaneous T-cell lymphoma (CTCL), large granular lymphocytic leukemia (LGF), Hodgkin's disease and Reed-Sternberg disease.

[365] Aberrant expression and/or activity of 50365 molecules may mediate disorders associated with bone metabolism. "Bone metabolism" refers to direct or indirect effects in the formation or degeneration of bone structures, e.g., bone formation, bone resorption, etc., which may ultimately affect the concentrations in serum of calcium and phosphate. This term also includes activities mediated by 50365 molecules effects in bone cells, e.g. osteoclasts and osteoblasts, that may in turn result in bone formation and degeneration. For example, 50365 molecules may support different activities of bone resorbing osteoclasts such as the stimulation of differentiation of monocytes and mononuclear phagocytes into osteoclasts. Accordingly, 50365 molecules that modulate the

production of bone cells can influence bone formation and degeneration, and thus may be used to treat bone disorders. Examples of such disorders include, but are not limited to, osteoporosis, osteodystrophy, osteomalacia, rickets, osteitis fibrosa cystica, renal osteodystrophy, osteosclerosis, anti-convulsant treatment, osteopenia, fibrogenesis-imperfecta ossium, secondary hyperparathyroidism, hypoparathyroidism, hyperparathyroidism, cirrhosis, obstructive jaundice, drug induced metabolism, medullary carcinoma, chronic renal disease, rickets, sarcoidosis, glucocorticoid antagonism, malabsorption syndrome, steatorrhea, tropical sprue, idiopathic hypercalcemia and milk fever.

10 [366] The 50365 nucleic acid and protein of the invention can be used to treat and/or diagnose a variety of immune disorders. Examples of immune disorders or diseases include, but are not limited to, autoimmune diseases (including, for example, diabetes mellitus, arthritis (including rheumatoid arthritis, juvenile rheumatoid arthritis, osteoarthritis, psoriatic arthritis), multiple sclerosis, encephalomyelitis, myasthenia gravis, 15 systemic lupus erythematosus, autoimmune thyroiditis, dermatitis (including atopic dermatitis and eczematous dermatitis), psoriasis, Sjögren's Syndrome, Crohn's disease, aphthous ulcer, iritis, conjunctivitis, keratoconjunctivitis, ulcerative colitis, asthma, allergic asthma, cutaneous lupus erythematosus, scleroderma, vaginitis, proctitis, drug eruptions, leprosy reversal reactions, erythema nodosum leprosum, autoimmune uveitis, allergic 20 encephalomyelitis, acute necrotizing hemorrhagic encephalopathy, idiopathic bilateral progressive sensorineural hearing loss, aplastic anemia, pure red cell anemia, idiopathic thrombocytopenia, polychondritis, Wegener's granulomatosis, chronic active hepatitis, Stevens-Johnson syndrome, idiopathic sprue, lichen planus, Graves' disease, sarcoidosis, primary biliary cirrhosis, uveitis posterior, and interstitial lung fibrosis), graft-versus-host 25 disease, cases of transplantation, and allergy such as, atopic allergy.

[367] Examples of disorders involving the heart or "cardiovascular disorder" include, but are not limited to, a disease, disorder, or state involving the cardiovascular system, *e.g.*, the heart, the blood vessels, and/or the blood. A cardiovascular disorder can be caused by an imbalance in arterial pressure, a malfunction of the heart, or an occlusion of a 30 blood vessel, *e.g.*, by a thrombus. Examples of such disorders include hypertension, atherosclerosis, coronary artery spasm, congestive heart failure, coronary artery disease, valvular disease, arrhythmias, and cardiomyopathies.

[368] Additionally, 50365 may play an important role in the regulation of metabolism or pain disorders. Diseases of metabolic imbalance include, but are not limited to, obesity, anorexia nervosa, cachexia, lipid disorders, and diabetes. Examples of pain disorders include, but are not limited to, pain response elicited during various forms of tissue injury, e.g., inflammation, infection, and ischemia, usually referred to as hyperalgesia (described in, for example, Fields, H.L. (1987) *Pain*, New York:McGraw-Hill); pain associated with musculoskeletal disorders, e.g., joint pain; tooth pain; headaches; pain associated with surgery; pain related to irritable bowel syndrome; or chest pain.

[369] As discussed, successful treatment of 50365 disorders can be brought about by techniques that serve to inhibit the expression or activity of target gene products. For example, compounds, e.g., an agent identified using an assays described above, that proves to exhibit negative modulatory activity, can be used in accordance with the invention to prevent and/or ameliorate symptoms of 50365 disorders. Such molecules can include, but are not limited to peptides, phosphopeptides, small organic or inorganic molecules, or antibodies (including, for example, polyclonal, monoclonal, humanized, anti-idiotypic, chimeric or single chain antibodies, and Fab, F(ab')₂ and Fab expression library fragments, scFV molecules, and epitope-binding fragments thereof).

[370] Further, antisense and ribozyme molecules that inhibit expression of the target gene can also be used in accordance with the invention to reduce the level of target gene expression, thus effectively reducing the level of target gene activity. Still further, triple helix molecules can be utilized in reducing the level of target gene activity. Antisense, ribozyme and triple helix molecules are discussed above.

[371] It is possible that the use of antisense, ribozyme, and/or triple helix molecules to reduce or inhibit mutant gene expression can also reduce or inhibit the transcription (triple helix) and/or translation (antisense, ribozyme) of mRNA produced by normal target gene alleles, such that the concentration of normal target gene product present can be lower than is necessary for a normal phenotype. In such cases, nucleic acid molecules that encode and express target gene polypeptides exhibiting normal target gene activity can be introduced into cells via gene therapy method. Alternatively, in instances in that the target gene encodes an extracellular protein, it can be preferable to co-administer normal target gene protein into the cell or tissue in order to maintain the requisite level of cellular or tissue target gene activity.

[372] Another method by which nucleic acid molecules may be utilized in treating or preventing a disease characterized by 50365 expression is through the use of aptamer molecules specific for 50365 protein. Aptamers are nucleic acid molecules having a tertiary structure which permits them to specifically bind to protein ligands (see, e.g., Osborne, *et al.* 5 (1997) *Curr. Opin. Chem Biol.* 1: 5-9; and Patel, D.J. (1997) *Curr Opin Chem Biol* 1:32-46). Since nucleic acid molecules may in many cases be more conveniently introduced into target cells than therapeutic protein molecules may be, aptamers offer a method by which 50365 protein activity may be specifically decreased without the introduction of drugs or other molecules which may have pluripotent effects.

10 [373] Antibodies can be generated that are both specific for target gene product and that reduce target gene product activity. Such antibodies may, therefore, be administered in instances whereby negative modulatory techniques are appropriate for the treatment of 50365 disorders. For a description of antibodies, see the Antibody section above.

[374] In circumstances wherein injection of an animal or a human subject with a 15 50365 protein or epitope for stimulating antibody production is harmful to the subject, it is possible to generate an immune response against 50365 through the use of anti-idiotypic antibodies (see, for example, Herlyn, D. (1999) *Ann Med* 31:66-78; and Bhattacharya-Chatterjee, M., and Foon, K.A. (1998) *Cancer Treat Res.* 94:51-68). If an anti-idiotypic antibody is introduced into a mammal or human subject, it should stimulate the production 20 of anti-anti-idiotypic antibodies, which should be specific to the 50365 protein. Vaccines directed to a disease characterized by 50365 expression may also be generated in this fashion.

[375] In instances where the target antigen is intracellular and whole antibodies are used, internalizing antibodies may be preferred. Lipofectin or liposomes can be used to 25 deliver the antibody or a fragment of the Fab region that binds to the target antigen into cells. Where fragments of the antibody are used, the smallest inhibitory fragment that binds to the target antigen is preferred. For example, peptides having an amino acid sequence corresponding to the Fv region of the antibody can be used. Alternatively, single chain neutralizing antibodies that bind to intracellular target antigens can also be administered. 30 Such single chain antibodies can be administered, for example, by expressing nucleotide sequences encoding single-chain antibodies within the target cell population (see e.g., Marasco *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:7889-7893).

[376] The identified compounds that inhibit target gene expression, synthesis and/or activity can be administered to a patient at therapeutically effective doses to prevent, treat or ameliorate 50365 disorders. A therapeutically effective dose refers to that amount of the compound sufficient to result in amelioration of symptoms of the disorders. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures as described above.

[377] The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED_{50} with little or no toxicity. The dosage can vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose can be formulated in animal models to achieve a circulating plasma concentration range that includes the IC_{50} (i.e., the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma can be measured, for example, by high performance liquid chromatography.

[378] Another example of determination of effective dose for an individual is the ability to directly assay levels of "free" and "bound" compound in the serum of the test subject. Such assays may utilize antibody mimics and/or "biosensors" that have been created through molecular imprinting techniques. The compound which is able to modulate 50365 activity is used as a template, or "imprinting molecule", to spatially organize polymerizable monomers prior to their polymerization with catalytic reagents. The subsequent removal of the imprinted molecule leaves a polymer matrix which contains a repeated "negative image" of the compound and is able to selectively rebind the molecule under biological assay conditions. A detailed review of this technique can be seen in Ansell, R. J. *et al* (1996) *Current Opinion in Biotechnology* 7:89-94 and in Shea, K.J. (1994) *Trends in Polymer Science* 2:166-173. Such "imprinted" affinity matrixes are amenable to ligand-binding assays, whereby the immobilized monoclonal antibody component is replaced by an appropriately imprinted matrix. An example of the use of such matrixes in this way can be seen in Vlatakis, G. *et al* (1993) *Nature* 361:645-647. Through the use of isotope-labeling, the "free" concentration of compound which modulates the expression or activity of 50365 can be readily monitored and used in calculations of IC_{50} .

[379] Such "imprinted" affinity matrixes can also be designed to include fluorescent groups whose photon-emitting properties measurably change upon local and selective binding of target compound. These changes can be readily assayed in real time using appropriate fiberoptic devices, in turn allowing the dose in a test subject to be quickly optimized based on its individual IC_{50} . An rudimentary example of such a "biosensor" is discussed in Kriz, D. *et al* (1995) *Analytical Chemistry* 67:2142-2144.

[380] Another aspect of the invention pertains to methods of modulating 50365 expression or activity for therapeutic purposes. Accordingly, in an exemplary embodiment, the modulatory method of the invention involves contacting a cell with a 50365 or agent that modulates one or more of the activities of 50365 protein activity associated with the cell. An agent that modulates 50365 protein activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring target molecule of a 50365 protein (e.g., a 50365 substrate or receptor), a 50365 antibody, a 50365 agonist or antagonist, a peptidomimetic of a 50365 agonist or antagonist, or other small molecule.

[381] In one embodiment, the agent stimulates one or 50365 activities. Examples of such stimulatory agents include active 50365 protein and a nucleic acid molecule encoding 50365. In another embodiment, the agent inhibits one or more 50365 activities. Examples of such inhibitory agents include antisense 50365 nucleic acid molecules, anti-50365 antibodies, and 50365 inhibitors. These modulatory methods can be performed *in vitro* (e.g., by culturing the cell with the agent) or, alternatively, *in vivo* (e.g., by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by aberrant or unwanted expression or activity of a 50365 protein or nucleic acid molecule. In one embodiment, the method involves administering an agent (e.g., an agent identified by a screening assay described herein), or combination of agents that modulates (e.g., up regulates or down regulates) 50365 expression or activity. In another embodiment, the method involves administering a 50365 protein or nucleic acid molecule as therapy to compensate for reduced, aberrant, or unwanted 50365 expression or activity.

[382] Stimulation of 50365 activity is desirable in situations in which 50365 is abnormally downregulated and/or in which increased 50365 activity is likely to have a beneficial effect. For example, stimulation of 50365 activity is desirable in situations in which a 50365 is downregulated and/or in which increased 50365 activity is likely to have a beneficial effect. Likewise, inhibition of 50365 activity is desirable in situations in which

50365 is abnormally upregulated and/or in which decreased 50365 activity is likely to have a beneficial effect.

Pharmacogenomics

- 5 [383] The 50365 molecules of the present invention, as well as agents, or modulators which have a stimulatory or inhibitory effect on 50365 activity (e.g., 50365 gene expression) as identified by a screening assay described herein can be administered to individuals to treat (prophylactically or therapeutically) 50365 associated disorders (e.g., a cell proliferative disorder or aberrant metabolic activity, e.g., aberrant phosphorylation of
- 10 aldo- and ketohexoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine)) associated with aberrant or unwanted 50365 activity. In conjunction with such treatment, pharmacogenomics (i.e., the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the
- 15 relation between dose and blood concentration of the pharmacologically active drug. Thus, a physician or clinician may consider applying knowledge obtained in relevant pharmacogenomics studies in determining whether to administer a 50365 molecule or 50365 modulator as well as tailoring the dosage and/or therapeutic regimen of treatment with a 50365 molecule or 50365 modulator.
- 20 [384] Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, for example, Eichelbaum, M. *et al.* (1996) *Clin. Exp. Pharmacol. Physiol.* 23:983-985 and Linder, M.W. *et al.* (1997) *Clin. Chem.* 43:254-266. In general, two types of pharmacogenetic conditions can be differentiated. Genetic conditions transmitted as a
- 25 single factor altering the way drugs act on the body (altered drug action) or genetic conditions transmitted as single factors altering the way the body acts on drugs (altered drug metabolism). These pharmacogenetic conditions can occur either as rare genetic defects or as naturally-occurring polymorphisms. For example, glucose-6-phosphate dehydrogenase deficiency (G6PD) is a common inherited enzymopathy in which the main clinical
- 30 complication is haemolysis after ingestion of oxidant drugs (anti-malarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

[385] One pharmacogenomics approach to identifying genes that predict drug response, known as "a genome-wide association", relies primarily on a high-resolution map of the human genome consisting of already known gene-related markers (e.g., a "bi-allelic" gene marker map which consists of 60,000-100,000 polymorphic or variable sites on the human genome, each of which has two variants.) Such a high-resolution genetic map can be compared to a map of the genome of each of a statistically significant number of patients taking part in a Phase II/III drug trial to identify markers associated with a particular observed drug response or side effect. Alternatively, such a high resolution map can be generated from a combination of some ten-million known single nucleotide polymorphisms (SNPs) in the human genome. As used herein, a "SNP" is a common alteration that occurs in a single nucleotide base in a stretch of DNA. For example, a SNP may occur once per every 1000 bases of DNA. A SNP may be involved in a disease process, however, the vast majority may not be disease-associated. Given a genetic map based on the occurrence of such SNPs, individuals can be grouped into genetic categories depending on a particular pattern of SNPs in their individual genome. In such a manner, treatment regimens can be tailored to groups of genetically similar individuals, taking into account traits that may be common among such genetically similar individuals.

[386] Alternatively, a method termed the "candidate gene approach," can be utilized to identify genes that predict drug response. According to this method, if a gene that encodes a drug's target is known (e.g., a 50365 protein of the present invention), all common variants of that gene can be fairly easily identified in the population and it can be determined if having one version of the gene versus another is associated with a particular drug response.

[387] Alternatively, a method termed the "gene expression profiling," can be utilized to identify genes that predict drug response. For example, the gene expression of an animal dosed with a drug (e.g., a 50365 molecule or 50365 modulator of the present invention) can give an indication whether gene pathways related to toxicity have been turned on.

[388] Information generated from more than one of the above pharmacogenomics approaches can be used to determine appropriate dosage and treatment regimens for prophylactic or therapeutic treatment of an individual. This knowledge, when applied to dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when treating a subject with a 50365 molecule or

50365 modulator, such as a modulator identified by one of the exemplary screening assays described herein.

[389] The present invention further provides methods for identifying new agents, or combinations, that are based on identifying agents that modulate the activity of one or more of the gene products encoded by one or more of the 50365 genes of the present invention, wherein these products may be associated with resistance of the cells to a therapeutic agent. Specifically, the activity of the proteins encoded by the 50365 genes of the present invention can be used as a basis for identifying agents for overcoming agent resistance. By blocking the activity of one or more of the resistance proteins, target cells, e.g., human cells, will become sensitive to treatment with an agent that the unmodified target cells were resistant to.

[390] Monitoring the influence of agents (e.g., drugs) on the expression or activity of a 50365 protein can be applied in clinical trials. For example, the effectiveness of an agent determined by a screening assay as described herein to increase 50365 gene expression, protein levels, or upregulate 50365 activity, can be monitored in clinical trials of subjects exhibiting decreased 50365 gene expression, protein levels, or downregulated 50365 activity. Alternatively, the effectiveness of an agent determined by a screening assay to decrease 50365 gene expression, protein levels, or downregulate 50365 activity, can be monitored in clinical trials of subjects exhibiting increased 50365 gene expression, protein levels, or upregulated 50365 activity. In such clinical trials, the expression or activity of a 50365 gene, and preferably, other genes that have been implicated in, for example, a 50365-associated disorder can be used as a "read out" or markers of the phenotype of a particular cell.

50365 Informatics

[391] The sequence of a 50365 molecule is provided in a variety of media to facilitate use thereof. A sequence can be provided as a manufacture, other than an isolated nucleic acid or amino acid molecule, which contains a 50365. Such a manufacture can provide a nucleotide or amino acid sequence, e.g., an open reading frame, in a form which allows examination of the manufacture using means not directly applicable to examining the nucleotide or amino acid sequences, or a subset thereof, as they exists in nature or in purified form. The sequence information can include, but is not limited to, 50365 full-length nucleotide and/or amino acid sequences, partial nucleotide and/or amino acid sequences,

polymorphic sequences including single nucleotide polymorphisms (SNPs), epitope sequence, and the like. In a preferred embodiment, the manufacture is a machine-readable medium, e.g., a magnetic, optical, chemical or mechanical information storage device.

[392] As used herein, "machine-readable media" refers to any medium that can be read and accessed directly by a machine, e.g., a digital computer or analogue computer. Non-limiting examples of a computer include a desktop PC, laptop, mainframe, server (e.g., a web server, network server, or server farm), handheld digital assistant, pager, mobile telephone, and the like. The computer can be stand-alone or connected to a communications network, e.g., a local area network (such as a VPN or intranet), a wide area network (e.g., an Extranet or the Internet), or a telephone network (e.g., a wireless, DSL, or ISDN network). Machine-readable media include, but are not limited to: magnetic storage media, such as floppy discs, hard disc storage medium, and magnetic tape; optical storage media such as CD-ROM; electrical storage media such as RAM, ROM, EPROM, EEPROM, flash memory, and the like; and hybrids of these categories such as magnetic/optical storage media.

[393] A variety of data storage structures are available to a skilled artisan for creating a machine-readable medium having recorded thereon a nucleotide or amino acid sequence of the present invention. The choice of the data storage structure will generally be based on the means chosen to access the stored information. In addition, a variety of data processor programs and formats can be used to store the nucleotide sequence information of the present invention on computer readable medium. The sequence information can be represented in a word processing text file, formatted in commercially-available software such as WordPerfect and Microsoft Word, or represented in the form of an ASCII file, stored in a database application, such as DB2, Sybase, Oracle, or the like. The skilled artisan can readily adapt any number of data processor structuring formats (e.g., text file or database) in order to obtain computer readable medium having recorded thereon the nucleotide sequence information of the present invention.

[394] In a preferred embodiment, the sequence information is stored in a relational database (such as Sybase or Oracle). The database can have a first table for storing sequence (nucleic acid and/or amino acid sequence) information. The sequence information can be stored in one field (e.g., a first column) of a table row and an identifier for the sequence can be stored in another field (e.g., a second column) of the table row. The database can have a second table, e.g., storing annotations. The second table can have a field for the

sequence identifier, a field for a descriptor or annotation text (e.g., the descriptor can refer to a functionality of the sequence, a field for the initial position in the sequence to which the annotation refers, and a field for the ultimate position in the sequence to which the annotation refers. Non-limiting examples for annotation to nucleic acid sequences include polymorphisms (e.g., SNP's) translational regulatory sites and splice junctions. Non-limiting examples for annotations to amino acid sequence include polypeptide domains, e.g., a domain described herein; active sites and other functional amino acids; and modification sites.

[395] By providing the nucleotide or amino acid sequences of the invention in computer readable form, the skilled artisan can routinely access the sequence information for a variety of purposes. For example, one skilled in the art can use the nucleotide or amino acid sequences of the invention in computer readable form to compare a target sequence or target structural motif with the sequence information stored within the data storage means. A search is used to identify fragments or regions of the sequences of the invention which match a particular target sequence or target motif. The search can be a BLAST search or other routine sequence comparison, e.g., a search described herein.

[396] Thus, in one aspect, the invention features a method of analyzing 50365, e.g., analyzing structure, function, or relatedness to one or more other nucleic acid or amino acid sequences. The method includes: providing a 50365 nucleic acid or amino acid sequence; comparing the 50365 sequence with a second sequence, e.g., one or more preferably a plurality of sequences from a collection of sequences, e.g., a nucleic acid or protein sequence database to thereby analyze 50365. The method can be performed in a machine, e.g., a computer, or manually by a skilled artisan.

[397] The method can include evaluating the sequence identity between a 50365 sequence and a database sequence. The method can be performed by accessing the database at a second site, e.g., over the Internet.

[398] As used herein, a "target sequence" can be any DNA or amino acid sequence of six or more nucleotides or two or more amino acids. A skilled artisan can readily recognize that the longer a target sequence is, the less likely a target sequence will be present as a random occurrence in the database. Typical sequence lengths of a target sequence are from about 10 to 100 amino acids or from about 30 to 300 nucleotide residues. However, it is well recognized that commercially important fragments, such as sequence fragments involved in gene expression and protein processing, may be of shorter length.

[399] Computer software is publicly available which allows a skilled artisan to access sequence information provided in a computer readable medium for analysis and comparison to other sequences. A variety of known algorithms are disclosed publicly and a variety of commercially available software for conducting search means are and can be used in the computer-based systems of the present invention. Examples of such software include, but are not limited to, MacPattern (EMBL), BLASTN and BLASTX (NCBI).

[400] Thus, the invention features a method of making a computer readable record of a sequence of a 50365 sequence which includes recording the sequence on a computer readable matrix. In a preferred embodiment the record includes one or more of the following: identification of an ORF; identification of a domain, region, or site; identification of the start of transcription; identification of the transcription terminator; the full length amino acid sequence of the protein, or a mature form thereof; the 5' end of the translated region.

[401] In another aspect, the invention features, a method of analyzing a sequence. The method includes: providing a 50365 sequence, or record, in machine-readable form; comparing a second sequence to the 50365 sequence; thereby analyzing a sequence. Comparison can include comparing to sequences for sequence identity or determining if one sequence is included within the other, e.g., determining if the 50365 sequence includes a sequence being compared. In a preferred embodiment the 50365 or second sequence is stored on a first computer, e.g., at a first site and the comparison is performed, read, or recorded on a second computer, e.g., at a second site. E.g., the 50365 or second sequence can be stored in a public or proprietary database in one computer, and the results of the comparison performed, read, or recorded on a second computer. In a preferred embodiment the record includes one or more of the following: identification of an ORF; identification of a domain, region, or site; identification of the start of transcription; identification of the transcription terminator; the full length amino acid sequence of the protein, or a mature form thereof; the 5' end of the translated region.

[402] In another aspect, the invention provides a machine-readable medium for holding instructions for performing a method for determining whether a subject has a 50365-associated disease or disorder or a pre-disposition to a 50365-associated disease or disorder, wherein the method comprises the steps of determining 50365 sequence information associated with the subject and based on the 50365 sequence information, determining whether the subject has a 50365-associated disease or disorder or a pre-disposition to a

50365-associated disease or disorder and/or recommending a particular treatment for the disease, disorder or pre-disease condition.

[403] The invention further provides in an electronic system and/or in a network, a method for determining whether a subject has a 50365-associated disease or disorder or a pre-disposition to a disease associated with a 50365 wherein the method comprises the steps of determining 50365 sequence information associated with the subject, and based on the 50365 sequence information, determining whether the subject has a 50365-associated disease or disorder or a pre-disposition to a 50365-associated disease or disorder, and/or recommending a particular treatment for the disease, disorder or pre-disease condition. In a preferred embodiment, the method further includes the step of receiving information, e.g., phenotypic or genotypic information, associated with the subject and/or acquiring from a network phenotypic information associated with the subject. The information can be stored in a database, e.g., a relational database. In another embodiment, the method further includes accessing the database, e.g., for records relating to other subjects, comparing the 50365 sequence of the subject to the 50365 sequences in the database to thereby determine whether the subject as a 50365-associated disease or disorder, or a pre-disposition for such.

[404] The present invention also provides in a network, a method for determining whether a subject has a 50365 associated disease or disorder or a pre-disposition to a 50365-associated disease or disorder associated with 50365, said method comprising the steps of receiving 50365 sequence information from the subject and/or information related thereto, receiving phenotypic information associated with the subject, acquiring information from the network corresponding to 50365 and/or corresponding to a 50365-associated disease or disorder (e.g., faulty glycolytic activity, including faulty phosphorylation of aldo- and ketohexoses (e.g., glucose, mannose, fructose, sorbitol and glucosamine)), and based on one or more of the phenotypic information, the 50365 information (e.g., sequence information and/or information related thereto), and the acquired information, determining whether the subject has a 50365-associated disease or disorder or a pre-disposition to a 50365-associated disease or disorder. The method may further comprise the step of recommending a particular treatment for the disease, disorder or pre-disease condition.

[405] The present invention also provides a method for determining whether a subject has a 50365 -associated disease or disorder or a pre-disposition to a 50365-associated disease or disorder, said method comprising the steps of receiving information related to 50365 (e.g., sequence information and/or information related thereto), receiving

phenotypic information associated with the subject, acquiring information from the network related to 50365 and/or related to a 50365-associated disease or disorder, and based on one or more of the phenotypic information, the 50365 information, and the acquired information, determining whether the subject has a 50365-associated disease or disorder or a pre-disposition to a 50365-associated disease or disorder. The method may further comprise the step of recommending a particular treatment for the disease, disorder or pre-disease condition.

[406] This invention is further illustrated by the following examples that should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are incorporated herein by reference.

EXAMPLES

Example 1: Identification and Characterization of Human 50365 cDNA

[407] The human 50365 nucleic acid sequence is recited as follows:

[408] CCACGCGTCCGGCCTGGACTGGAAGCGTGCAAACTCCAGAGTCG
15 TAGGAGTGAACACTGCACAGGAATCTCTGCCCATCTCAGGAGAAACCAAACCTTG
GGGAAAATGTTTGC GGTCCTTGTATGGCATTCTTACTTCAGCAAGCTGAAGGAG
GACCAGATCAAGAAGGTGGACAGGTTCTGTATCACATGCGGCTCTCCGATGAC
ACCCTTTTGGACATCATGAGGCGGTTCCGGGCTGAGATGGAGAAGGGCCTGGCA
AAGGACACCAACCCACGGCTGCAGTGAAGATGTTGCCACCTTCGTCAGGGCC
20 ATTCCCGATGGTTCCGAAAATGGGGAGTTCCTTTCCCTGGATCTCGGAGGGTCC
AAGTTCCGAGTGCTGAAGGTGCAAGTCGCTGAAGAGGGGAAGCGACACGTGCA
GATGGAGAGTCAGTTCTACCCAACGCCCAATGAAATCATCCGCGGGAACGGCAT
AGAGCTGTTTGAATATGTAGCTGACTGTCTGGCAGATTTTCATGAAGACCAAAGA
TTTAAAGCATAAGAAATTGCCCTTGGCCTAACTTTTTCTTTCCCTGTCTGACAG
25 ACTAAACTGGAAGAGGGTGTCTACTTTCGTGGACAAAAAAGTTTAAGGCACGA
GGAGTTCAGGACACGGATGTGGTGAGCCGTCTGACCAAAGCCATGAGAAGACA
CAAGGACATGGACGTGGACATCCTGGCCCTGGTCAATGACACCGTGGGGACCAT
GATGACCTGTGCCTATGACGACCCCTACTGCGAAGTTGGTGTTCATCATCGGAAC
TGGCACCAATGCGTGTTACATGGAGGACATGAGCAACATTGACCTGGTGGAGG
30 GCGACGAGGGCAGGATGTGCATCAACACAGAGTGGGGGGCCTTCGGGGACGAC

GGGGCCCTGGAGGACATTCGCACTGAGTTCGACAGGGAGCTGGACCTCGGCTCT
CTCAACCCAGGAAAGCAACTGTTGAGAAGATGATCAGTGGCCTGTACCTGGGG
GAGCTTGTGAGGCTTATCTTGCTGAAGATGGCCAAGGCTGGCCTCCTGTTTGGTG
GTGAGAAATCTTCTGCTCTCCACACTAAGGGCAAGATCGAAACACGGCACGTGG
5 CTGCCATGGAGAAGTATAAAGAAGGCCTTGCTAATAACAAGAGAGATCCTGGTG
GACCTGGGTCTGGAACCGTCTGAGGCTGACTGCATTGCCGTCCAGCATGTCTGT
ACCATCGTCTCCTTCCGCTCGGCCAATCTCTGTGCAGCAGCTCTGGCGGCCATCC
TGACACGCCTCCGGGAGAACAAGAAGGTGGAACGGCTCCGGACCACAGTGGGC
ATGGACGGCACCCCTCTACAAGATACACCCTCAGTACCCAAAACGCCTGCACAAG
10 GTGGTGAGGAACTGGTCCCAAGCTGTGATGTCCGCTTCTCCTGTGAGAGAGT
GGCAGCACCAAGGGGGCCGCCATGGTGACCGCGGTGGCCTCCCGCGTGCAGGC
CCAGCGGAAGCAGATCGACAGGGTGCTGGCTTTGTTCCAGCTGACCCGAGAGCA
GCTCGTGGACGTGCAGGCCAAGATGCGGGCTGAGCTGGAGTATGGGCTGAAGA
AGAAGAGCCACGGGCTGGCCACGGTCAGGATGCTGCCCACCTACGTCTGCGGG
15 CTGCCGGACGGCACAGAGAAAGGAAAGTTTCTCGCCCTGGATCTTGGGGGAAC
CAACTTCCGGGTCTCCTGGTGAAGATCAGAAAGTGGACGGAGGTCAGTGCGAAT
GTACAACAAGATCTTCGCCATCCCCCTGGAGATCATGCAGGGCACTGGTGAGGA
GCTCTTTGATCACATTGTGCAGTGCATCGCCGACTTCCTGGACTACATGGGCCTC
AAGGGAGCCTCCCTACCTTTGGGCTTCACATTCTCATTTCCCTGCAGGCAGATGA
20 GCATTGACAAGGGAACACTCATAGGGTGGACCAAAGGTTTCAAGGCCACTGAC
TGTGAAGGGGAGGACGTGGTGGACATGCTCAGGGAAGCCATCAAGAGGAGAAA
CGAGTTTGACCTGGACATTGTTGCAGTCGTGAATGATACAGTGGGGACCATGAT
GACCTGTGGCTATGAAGATCCTAATTGTGAGATTGGCCTGATTGCAGGAACAGG
CAGCAACATGTGCTACATGGAGGACATGAGGAACATCGAGATGGTGGAGGGGG
25 GTGAAGGGAAGATGTGCATCAATACAGAGTGGGGAGGATTTGGAGACAATGGC
TGCATAGATGACATCCGGACCCGATACGACACGGAGGTGGATGAGGGGTCTT
GAATCCTGGCAAGCAGAGATACGAGAAAATGACCAGTGGGATGTACTTGGGGG
AGATTGTGCGGCAGATCCTGATCGACCTGACCAAGCAGGGTCTCCTCTTCCGAG
GGCAGATTTTCAAGCGTCTCCGGACCAGGGGCATCTTCGAAACCAAGTTCCTGT
30 CCCAGATCGAAAGCGATCGGCTGGCCCTTCTCCAGGTGAGGAGGATTCTGCAGC
AGCTGGGCCTGGACAGCACGTGTGAGGACAGCATCGTGGTGAAGGAGGTGTGC
GGAGCCGTGTCCCGGCGGGCGGCCAGCTCTGCGGTGCTGGCCTGGCCGCTATA
GTGGAAAAAAGGAGAGAAGACCAGGGGCTAGAGCACCTGAGGATCACTGTGGG

TGTGGACGGCACCCCTGTACAAGCTGCACCCTCACTTTTCTAGAATATTGCAGGA
AACTGTGAAGGAACTAGCCCCCTCGATGTGATGTGACATTCATGCTGTCAGAAGA
TGGCAGTGGAAAAGGGGCAGCACTGATCACTGCTGTGGCCAAGAGGTTACAGC
AGGCACAGAAGGAGAACT**AGGA**ACCCCTGGGATTGGACCTGATGCATCTTGG
5 TACTGAACAGCTTTTCTCTGGCAGATCAGTTGGTCAGAGACCAATGGGCACCC
TCCTGGCTGACCTCACCTTCTGGATGGCCGAAAGAGAACCCAGGTTCTCGGGT
ACTCTTAGTATCTTGTACTGGATTTCAGTGACATTACATGACATCTCTATTTGG
TATATTTGGGCCAAAATGGGCCAACTTATGAAATCAAAGTGTCTGTCCTGAGAG
ATCCCCCTTCAACACATTGTTTCAGGTGAGGCTTGAGCTGTCAATTCTCTATGGCT
10 TTCAGTCTTGTGGCTGCGGGACTTGGAATATATAGAATCTGCCCATGTGGCTG
GCAGGCTGTTTCCCCATTGGGATGCTTAAGCCATCTCTTATAGGGGATTGGACCC
TGTACTTGTGGATGAACATTGGAGAGCAAGAGGAACTCACGTTATGAACTAGGG
GGATCTCATCTAACTTGTCTTAAGTTGCCATGTTGACTTCAAACCTGTTAAGAG
AACAAAGACTTTGAAGTATCCAGCCCCAGGGTGCAGAGAGGTTGATTGCCAGG
15 GAGCACTGCAGGAATCATTGCATGCTTAAAGCGAGTTATGTCAGCACCCCTGTAG
GATTTTGTTCCTTATTAAGTGTGTGCCATGTGGTGGGGTGCTGTCTGGGGCATCT
GTTTTTCATTTTGCCTGTGGTTTGTGTTGCAGSTGTTGATAGTTGTTTTAAGGATT
GTTAGGTATAGGAAATCCAGTAAATTAATAAAAAAATTTTGATTTTCCAATAAA
AAAAAAAAAAAAAAAA (SEQ ID NO:1).

20 [409] The human 50365 sequence (SEQ ID NO:1) is approximately 3669
nucleotides long. The nucleic acid sequence includes an initiation codon (ATG) and a
termination codon (TAA) which are underscored and bolded above. The region between
and inclusive of the initiation codon and the termination codon is a methionine-initiated
coding sequence of about 2754 nucleotides (SEQ ID NO:3), including the termination
25 codon. The coding sequence encodes a 917 amino acid protein (SEQ ID NO:2), which is
recited as follows:

[410] MFAVHLMAFYFSKLKEDQIKKVDRFLYHMRLSDDTLLDIMRRFRAE
MEKGLAKDTNPTAAVKMLPTFVRAIPDGSENGEFLSLDLGGSKFRVLKVQVAEEG
KRHVQMESQFYPTPNEIIRNGNIELFYVADCLADFMKTKDLKHKKLPLGLTFSFPC
30 RQTKLEEGVLLSWTKKFKARGVQDQDVVSRLTKAMRRHKDMDVDILALVNDTVG
TMMTCAYDDPYCEVGVIIGTGTNACYMEDMSNIDLVEGDEGRMCINTEWGAFGD
DGALEDIRTEFDRELDLGSNPGKQLFEKMISGLYLGELVRLILLKMAKAGLLFGGE
KSSALHTKGKIETRHVAAMEKYKEGLANTREILVDLGLPSEADCIQHVCTIVSF

RSANLCAAALAILTRLRENKKVERLRTTVGMDGTLYKIHPQYPKRLHKVVRKLV
 SCDVRFLLESSTKGAAMVTAVASRVQAQRKQIDRVLALFQLTREQLVDVQAKM
 RAELEYGLKKKSHGLATVRMLPTYVCGLPDGTEKGKFLALDLGGTNFRVLLVKIRS
 GRRSVRMYNKIFAIPLEIMQGTGEELFDHIVQCIADFLDYMGLKGASLPLGFTSFPC
 5 RQMSIDKGTLLIGWTKGFKATDCEGEDVVDMLREAIKRRNEFDLDIVAVVNDTVGT
 MMTTCGYEDPNCEIGLIAGTGSNMCYMEDMRNIEMVEGGEGKMCINTEWGGFGDN
 GCIDDIRTRYDTEVDEGSLNPGKQRYEKMTSGMYLGEIVRQILIDLTKQGLLFRGQIS
 ERLRTRGIFETKFLSQIESDRLALLQVRRILQQLGLDSTCEDSVVKEVCGAVSRAA
 QLCGAGLAAIVEKRREDQGLEHLRITVGVDGTLYKLHPHFSRILQETVKELAPRCDV
 10 TFMLSEDGSGKGAALITAVAKRLQQAQKEN (SEQ ID NO:2).

Example 2: Tissue Distribution of 50365 mRNA by TaqMan Analysis

[411] Endogenous human 50365 gene expression was determined using the Perkin-
 Elmer/ABI 7700 Sequence Detection System which employs TaqMan technology. Briefly,
 TaqMan technology relies on standard RT-PCR with the addition of a third gene-specific
 15 oligonucleotide (referred to as a probe) which has a fluorescent dye coupled to its 5' end
 (typically 6-FAM) and a quenching dye at the 3' end (typically TAMRA). When the
 fluorescently tagged oligonucleotide is intact, the fluorescent signal from the 5' dye is
 quenched. As PCR proceeds, the 5' to 3' nucleolytic activity of Taq polymerase digests the
 labeled primer, producing a free nucleotide labeled with 6-FAM, which is now detected as a
 20 fluorescent signal. The PCR cycle where fluorescence is first released and detected is
 directly proportional to the starting amount of the gene of interest in the test sample, thus
 providing a quantitative measure of the initial template concentration. Samples can be
 internally controlled by the addition of a second set of primers/probe specific for a
 housekeeping gene such as GAPDH or β 2-microglobulin which has been labeled with a
 25 different fluorophore on the 5' end (typically VIC).

[412] To determine the level of 50365 in various human tissues a primer/probe set
 was designed. Total RNA was prepared from a series of human tissues using an RNeasy kit
 from Qiagen. First strand cDNA was prepared from 1 μ g total RNA using an oligo-dT
 primer and Superscript II reverse transcriptase (Gibco/BRL). cDNA obtained from
 30 approximately 50 ng total RNA was used per TaqMan reaction. Tissues tested include the
 human tissues and several cell lines shown in Tables 1 and 2 below. 50365 mRNA was
 detected in colon and liver tissue, and upregulated in colonic liver metastases (Table 1). In

addition, 50365 mRNA was also detectable in adenomas and adenocarcinomas. 50365 expression was also found to a lesser extent in some lung tumor and ovary tumor tissues (Table 2).

[413] Table 1

Tissue Type	Expression
CHT 410 Colon Normal	0.32
CHT 425 Colon Normal	0.41
CHT 371 Colon Normal	1.06
PIT 281 Colon Normal	0.00
NDR 211 Colon Normal	0.22
CHT 122 Adenomas	0.21
CHT 887 Adenomas	1.65
CHT 414 Colonic Adenocarcinoma-B	0.47
CHT 841 Colonic Adenocarcinoma-B	0.05
CHT 890 Colonic Adenocarcinoma-B	0.58
CHT 910 Colonic Adenocarcinoma-B	3.85
CHT 377 Colonic Adenocarcinoma-B	0.00
CHT 520 Colonic Adenocarcinoma-C	0.80
CHT 596 Colonic Adenocarcinoma-C	0.77
CHT 907 Colonic Adenocarcinoma-C	2.41
CHT 372 Colonic Adenocarcinoma-C	2.09
NDR 210 Colonic Adenocarcinoma-C	0.95
CHT 1365 Colonic Adenocarcinoma-C	2.54
CLN 740 Liver Normal	0.00
CLN 741 Liver Normal	0.00
NDR 165 Liver Normal	0.00
NDR 150 Liver Normal	0.14
PIT 236 Liver Normal	0.00
CHT 1878 Liver Normal	0.00
CHT 119 Colon Liver Metastasis	7.52
CHT 131 Colon Liver Metastasis	0.77
CHT 218 Colon Liver Metastasis	5.45
CHT 739 Colon Liver Metastasis	10.53
CHT 755 Colon Liver Metastasis	3.64
CHT 215 Colon Abdominal Metastasis	0.24
PIT 337 Colon Normal	0.29
CHT 807 Colonic Adenocarcinoma-B	61.64
CHT 382 Colonic Adenocarcinoma-B	57.11
CHT 077 Colon Liver Metastasis	180.49

The mRNA expression data for 50365 mRNA tabulated in Table 1 indicate that

5 50365 expression is upregulated in some adenomas and adenocarcinomas, and in most colonic liver metastases (see "Relative Expression" values). Relative expression in Table 1 is relative to expression of β 2-macroglobulin.

[414] Table 2

Tissue Type	Expression
PIT 400 Breast Normal	0.00
PIT 372 Breast Normal	0.00
CHT 559 Breast Normal	0.00
MDA 236-Breast Tumor:PD-IDC(ILC?)	0.00
MDA 304 Breast Tumor: MD-IDC	0.00
CHT 2002 Breast Tumor: IDC	0.00
CHT 562 Breast Tumor: IDC	0.00
NDR 138 Breast Tumor ILC (LG)	0.00
CHT 1841 Lymph node (Breast Metastasis)	0.00
PIT 58 Lung (Breast Metastasis)	0.00
CHT 620 Ovary Normal	0.00
PIT 208 Ovary Normal	0.00
CLN 012 Ovary Tumor	0.00
CLN 07 Ovary Tumor	0.05
CLN 17 Ovary Tumor	1.38
MDA 25 Ovary Tumor	0.00
MDA 216 Ovary Tumor	0.00
PIT 298 Lung Normal	0.00
MDA 185 Lung Normal	0.00
CLN 930 Lung Normal	0.00
MPI 215 Lung Tumor--SmC	0.00
MDA 259 Lung Tumor-PDNSCCL	0.00
CHT 832 Lung Tumor-PDNSCCL	0.97
MDA 262 Lung Tumor- Small Cell Carcinoma	0.00
CHT 793 Lung Tumor-Adenocarcinoma	0.03
CHT 331 Lung Tumor-Adenocarcinoma	0.00
CHT 405 Colon Normal	0.16
CHT 523 Colon Normal	0.65
CHT 371 Colon Normal	2.38
CHT 382 Colon Tumor: MD	0.88
CHT 528 Colon Tumor: MD	7.84
CLN 609 Colon Tumor	2.21
NDR 210 Colon Tumor: MD-PD	0.84
CHT 340 Colon-Liver Metastasis	3.23
NDR 100 Colon-Liver Metastasis	1.11
PIT 260 Liver Normal (female)	1.17
CHT 1653 Cervix Squamous CC	0.00
CHT 569 Cervix Squamous CC	0.00
A24 HMVEC-Arr	0.00
C48 HMVEC-Prol	0.00
Pooled Hemangiomas	0.00
HCT116N22 Normal Ox.ygenation	0.97
HCT116H22 Hypoxic	0.00

50365 mRNA was analyzed by TaqMan in a number of cell lines derived from normal and tumor cells (Table 2). Relative expression in Table 2 is relative to expression of β 2-macroglobulin. Elevated 50365 mRNA expression levels were detected in some colon cell lines, e.g., normal colon, colon tumor; colonic liver metastases; some lung cell lines, e.g., lung tumor-PDNSCCL (poorly differentiated non-small cell carcinoma of the lung), lung tumor-adenocarcinoma; and an ovary tumor cell line. 50365 mRNA was also detected under normal oxygenation conditions.

Example 3: Tissue Distribution of 50365 mRNA by Northern Analysis

[415] Northern blot hybridizations with various RNA samples can be performed under standard conditions and washed under stringent conditions, i.e., 0.2xSSC at 65°C. A DNA probe corresponding to all or a portion of the 50365 cDNA (SEQ ID NO:1) can be used. The DNA was radioactively labeled with ^{32}P -dCTP using the Prime-It Kit (Stratagene, La Jolla, CA) according to the instructions of the supplier. Filters containing mRNA from mouse hematopoietic and endocrine tissues, and cancer cell lines (Clontech, Palo Alto, CA) can be probed in ExpressHyb hybridization solution (Clontech) and washed at high stringency according to manufacturer's recommendations.

Example 4: Recombinant Expression of 50365 in Bacterial Cells

[416] In this example, 50365 is expressed as a recombinant glutathione-S-transferase (GST) fusion polypeptide in *E. coli* and the fusion polypeptide is isolated and characterized. Specifically, 50365 is fused to GST and this fusion polypeptide is expressed in *E. coli*, e.g., strain PEB199. Expression of the GST-50365 fusion protein in PEB199 is induced with IPTG. The recombinant fusion polypeptide is purified from crude bacterial lysates of the induced PEB199 strain by affinity chromatography on glutathione beads. Using polyacrylamide gel electrophoretic analysis of the polypeptide purified from the bacterial lysates, the molecular weight of the resultant fusion polypeptide is determined.

Example 5: Expression of Recombinant 50365 Protein in COS Cells

[417] To express the 50365 gene in COS cells, the pcDNA/Amp vector by Invitrogen Corporation (San Diego, CA) is used. This vector contains an SV40 origin of replication, an ampicillin resistance gene, an *E. coli* replication origin, a CMV promoter

- followed by a polylinker region, and an SV40 intron and polyadenylation site. A DNA fragment encoding the entire 50365 protein and an HA tag (Wilson *et al.* (1984) *Cell* 37:767) or a FLAG tag fused in-frame to its 3' end of the fragment is cloned into the polylinker region of the vector, thereby placing the expression of the recombinant protein under the control of the CMV promoter.
- 5 [418] To construct the plasmid, the 50365 DNA sequence is amplified by PCR using two primers. The 5' primer contains the restriction site of interest followed by approximately twenty nucleotides of the 50365 coding sequence starting from the initiation codon; the 3' end sequence contains complementary sequences to the other restriction site of interest, a translation stop codon, the HA tag or FLAG tag and the last 20 nucleotides of the 10 50365 coding sequence. The PCR amplified fragment and the pCDNA/Amp vector are digested with the appropriate restriction enzymes and the vector is dephosphorylated using the CIAP enzyme (New England Biolabs, Beverly, MA). Preferably the two restriction sites chosen are different so that the 50365 gene is inserted in the correct orientation. The 15 ligation mixture is transformed into *E. coli* cells (strains HB101, DH5 α , SURE, available from Stratagene Cloning Systems, La Jolla, CA, can be used), the transformed culture is plated on ampicillin media plates, and resistant colonies are selected. Plasmid DNA is isolated from transformants and examined by restriction analysis for the presence of the correct fragment.
- 20 [419] COS cells are subsequently transfected with the 50365-pcDNA/Amp plasmid DNA using the calcium phosphate or calcium chloride co-precipitation methods, DEAE-dextran-mediated transfection, lipofection, or electroporation. Other suitable methods for transfecting host cells can be found in Sambrook, J., Fritsh, E. F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. The expression of the 50365 25 polypeptide is detected by radiolabelling (^{35}S -methionine or ^{35}S -cysteine available from NEN, Boston, MA, can be used) and immunoprecipitation (Harlow, E. and Lane, D. (1988) *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY) using an HA specific monoclonal antibody. Briefly, the cells are labeled for 8 30 hours with ^{35}S -methionine (or ^{35}S -cysteine). The culture media are then collected and the cells are lysed using detergents (RIPA buffer, 150 mM NaCl, 1% NP-40, 0.1% SDS, 0.5% DOC, 50 mM Tris, pH 7.5). Both the cell lysate and the culture media are precipitated with

an HA specific monoclonal antibody. Precipitated polypeptides are then analyzed by SDS-PAGE.

- [420] Alternatively, DNA containing the 50365 coding sequence is cloned directly into the polylinker of the pCDNA/Amp vector using the appropriate restriction sites. The
- 5 resulting plasmid is transfected into COS cells in the manner described above, and the expression of the 50365 polypeptide is detected by radiolabelling and immunoprecipitation using a 50365 specific monoclonal antibody.

Equivalents

- [421] Those skilled in the art will recognize, or be able to ascertain using no more
- 10 than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

1 What is claimed is:

- 1 1. An isolated nucleic acid molecule selected from the group consisting of:
 - 2 a) a nucleic acid molecule comprising a nucleotide sequence which is at least
3 80% identical to the nucleotide sequence of SEQ ID NO:1, or SEQ ID NO:3;
 - 4 b) a nucleic acid molecule comprising a fragment of at least 300 nucleotides of
5 the nucleotide sequence of SEQ ID NO: 1, or SEQ ID NO:3;
 - 6 c) a nucleic acid molecule which encodes a polypeptide comprising the amino
7 acid sequence of SEQ ID NO:2;
 - 8 d) a nucleic acid molecule which encodes a fragment of a polypeptide
9 comprising the amino acid sequence of SEQ ID NO:2, wherein the fragment comprises at
10 least 15 contiguous amino acids of SEQ ID NO: 2; and
 - 11 e) a nucleic acid molecule which encodes a naturally occurring allelic variant of
12 a polypeptide comprising the amino acid sequence of SEQ ID NO:2, wherein the nucleic
13 acid molecule hybridizes to a nucleic acid molecule comprising SEQ ID NO: 1, 3, or a
14 complement thereof, under stringent conditions.
- 1 2. The isolated nucleic acid molecule of claim 1, which is selected from the
2 group consisting of:
 - 3 a) a nucleic acid comprising the nucleotide sequence of SEQ ID NO: 1, SEQ ID
4 NO:3; and
 - 5 b) a nucleic acid molecule which encodes a polypeptide comprising the amino
6 acid sequence of SEQ ID NO:2.
- 1 3. The nucleic acid molecule of claim 1 further comprising a vector nucleic acid
2 sequence.
- 1 4. The nucleic acid molecule of claim 1 further comprising a nucleic acid
2 sequence encoding a heterologous polypeptide.
- 1 5. A host cell which contains the nucleic acid molecule of claim 1.
- 1 6. The host cell of claim 5 which is a mammalian host cell.

1 7. A non-human mammalian host cell containing the nucleic acid molecule of
2 claim 1.

1 8. An isolated polypeptide selected from the group consisting of:

2 a) a polypeptide which is encoded by a nucleic acid molecule comprising a
3 nucleotide sequence which is at least 80% identical to a nucleic acid comprising the
4 nucleotide sequence of SEQ ID NO: 1 or SEQ ID NO:3;

5 b) a naturally occurring allelic variant of a polypeptide comprising the amino
6 acid sequence of SEQ ID NO:2, wherein the polypeptide is encoded by a nucleic acid
7 molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO: 1, SEQ ID
8 NO:3, or a complement thereof under stringent conditions; and

9 c) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID
10 NO:2, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2.

1 9. The isolated polypeptide of claim 8 comprising the amino acid sequence of
2 SEQ ID NO:2.

1 10. The polypeptide of claim 8 further comprising a heterologous amino acid
2 sequence.

1 11. An antibody which selectively binds to a polypeptide of claim 8.

1 12. A method for producing a polypeptide selected from the group consisting of:

2 a) a polypeptide comprising the amino acid sequence of SEQ ID NO:2;

3 b) a polypeptide comprising a fragment of the amino acid sequence of SEQ ID
4 NO:2, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2;
5 and

6 c) a naturally occurring allelic variant of a polypeptide comprising the amino
7 acid sequence of SEQ ID NO:2, wherein the polypeptide is encoded by a nucleic acid
8 molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID
9 NO:3, or a complement thereof under stringent conditions;

10 the method comprising culturing the host cell of claim 5 under conditions in which
11 the nucleic acid molecule is expressed.

1 13. A method for detecting the presence of a polypeptide of claim 8 in a sample,
2 comprising:

- 3 a) contacting the sample with a compound which selectively binds to a
4 polypeptide of claim 8; and
5 b) determining whether the compound binds to the polypeptide in the sample.

1 14. The method of claim 13, wherein the compound which binds to the
2 polypeptide is an antibody.

1 15. A kit comprising a compound which selectively binds to a polypeptide of
2 claim 8 and instructions for use.

1 16. A method for detecting the presence of a nucleic acid molecule of claim 1 in
2 a sample, comprising the steps of:

- 3 a) contacting the sample with a nucleic acid probe or primer which selectively
4 hybridizes to the nucleic acid molecule; and
5 b) determining whether the nucleic acid probe or primer binds to a nucleic acid
6 molecule in the sample.

1 17. The method of claim 16, wherein the sample comprises mRNA molecules
2 and is contacted with a nucleic acid probe.

1 18. A kit comprising a compound which selectively hybridizes to a nucleic acid
2 molecule of claim 1 and instructions for use.

1 19. A method for identifying a compound which binds to a polypeptide of claim
2 8 comprising the steps of:

- 3 a) contacting a polypeptide, or a cell expressing a polypeptide of claim 8 with a
4 test compound; and
5 b) determining whether the polypeptide binds to the test compound.

1 20. The method of claim 19, wherein the binding of the test compound to the
2 polypeptide is detected by a method selected from the group consisting of:

- 3 a) detection of binding by direct detecting of test compound/polypeptide
4 binding;
5 b) detection of binding using a competition binding assay;
6 c) detection of binding using an assay for 50365-mediated catalytic activity.

1 21. A method for modulating the activity of a polypeptide of claim 8 comprising
2 contacting a polypeptide or a cell expressing a polypeptide of claim 8 with a compound
3 which binds to the polypeptide in a sufficient concentration to modulate the activity of the
4 polypeptide.

1 22. A method for identifying a compound which modulates the activity of a
2 polypeptide of claim 8, comprising:
3 a) contacting a polypeptide of claim 8 with a test compound; and
4 b) determining the effect of the test compound on the activity of the polypeptide
5 to thereby identify a compound which modulates the activity of the polypeptide.

1 23. A method of inhibiting proliferation, or inducing the killing, of a 50365-
2 expressing hyperproliferative cell, comprising contacting the hyperproliferative cell with a
3 compound that modulates the activity or expression of a polypeptide of claim 8, in an
4 amount which is effective to reduce or inhibit the proliferation of, or induce the killing of,
5 the hyperproliferative cell.

1 24. The method of claim 23, wherein the compound is selected from the group
2 consisting of a peptide, a phosphopeptide, a small organic molecule, a small inorganic
3 molecule and an antibody.

1 25. The method of claim 23, wherein the compound is an antibody conjugated to
2 a therapeutic moiety selected from the group consisting of a cytotoxin, a cytotoxic agent and
3 a radioactive metal ion.

1 26. A method of inhibiting proliferation, or inducing the killing, of a 50365-
2 expressing hyperproliferative cell, comprising contacting the hyperproliferative cell with a
3 compound that modulates the activity or expression of a nucleic acid molecule of claim 1, in

4 an amount which is effective to reduce or inhibit the proliferation of, or induce the killing of,
5 the hyperproliferative cell.

1 27. The method of claim 26, wherein the compound is an antisense, a ribozyme,
2 or a triple helix molecule.

1 28. The method of claim 23, wherein the hyperproliferative cell is found in a
2 solid tumor, a soft tissue tumor, or a metastatic lesion.

1 29. The method of claim 28, wherein the hyperproliferative cell is found in a
2 cancer of the colon or a metastasis of colonic origin.

1/5




Figure 1

2/5

Hexokinase Domain 1 Alignment

hexokinase: domain 1/2, from 16 to 463: score 837.2, E=5.6e-248

```

HMM      *->adllqaveellddFtvstEtLrevtkrfikemekGLsPPkegntAs
          +d++++vv+ +l++++s++tL+++++rf+ emekGL+ k++++tA+
50365    16   EDQIKKVDRLFVHMRSLSDDTLLDIMRRFRAEMEKGLA--KDTNPATA 60

          vvkmlPtFvRstPtGtEkGdFLALDLGGTNfRVllVklgngkg.vemtq
          vkmlPtFvR++P+G+E G+FL+LDLGG+ fRVl V++ ++gk++v+m++
50365    61 -VKMLPTFVRAIPDGSENGFEFLSLDLGGSKFRVLKVQVAEEGKRhVQMES 109

          skYriPeelmtgenvtgeqLFDFiAecikdFmdeqfpgkkekLpLGFTF
          + Y +P e+ +g +g +lF+++A+c++dFm+ + +k+kk LpLG TF
50365    110 QFYPTPNEIIRG---NGIELFEYVADCLADFMKTKDLKHKK--LPLGLTF 154

          SFpCsQtsInegiLiRwTKGFkiGRATnsgvEGhDVVqLLreAikrrGaf
          SFpC+Qt++ eg+L++WTK Fk+ +gv DVV L++A +r +
50365    155 SFPCRQTKLEEGVLLSWTKKfKA-----RGVQDtdVVSRLTKAMRRHKDM 199

          pidVVAvvNDTVGTlmscaYtkGRGdpecetviGIvGTGtNaCYmEemr
          ++d+ A+vNDTVGT+m+cAY+ dp ce +G+I+GTGtNaCYmE m+
50365    200 DVDILALVNDTVGTMMTCAYD---DPYCE--VGVIIGTGTNACYMEDMS 243

```

Figure 2A

3/5

Hexokinase Domain 1 Alignment (cont'd)

```

nlekleGkLkddipdegrMcINmEWGaFGDngldlprTkYDVvviDeeSp
nI+++eG      degrMcIN+EWGaFGD+g+l+++rT+++D+++D +S+
50365 244 NIDLVEG-----DEGRMCINTEWGAFGDDGALEDIRTEFDRELDLGS 286

NPGqQLFEKMISGmYLGEivRlilldLtkeglLFkgqdsplkttrgsfet
NPG+QLFEKMISG+YLGE+vRlil+++k glLF+g+ s++l t g++eT
50365 287 NPGKQLFEKMISGLYLGEIVRLILLLKMAKAGLLFGGEKSSALHTKGKIET 336

svlSrIEsDpsenledvrailqtaLgletTdeerklvrrvCeaVstRAAr
+++++++E+  +e+l+++r+iL + Lgle+++ +++ v++vC++Vs+R+A+
50365 337 RHVAAMEKY-KEGLANTREILVD-LGLEPSEADCIHVCTIVSFRSAN 384

LcaaglAAilkkirenigrerlktvGVdGsvYklyPgFkerlaeLrdl
Lcaa+lAAil ++ren++ erl+++tVG DG++Yk +P++ +rl++ +r l
50365 385 LCAALAAIILRLRENKKVERLRTTVGMDGTLTKIHPQYPKRLHKVVRKL 434

lpdcegseedkksvipAEDGSGkGAAlvaAVAakl<-* (SEQ ID NO:4)
+p+c      d v+++ +E GS kGAA+v+AVA +
50365 435 VPSC-----D--VRFLSESGSTKGAAmVTAVASRV 463 (SEQ ID NO:2)

```

Figure 2B

4/5

Hexokinase Domain 2 Alignment

hexokinase: domain 2/2, from 464 to 910: score 955.2, E=1.7e-283
HMM *-->adllqaveellddftvstEtLrevtkrfikemekGLsppkeggntAs
+++ +++ l+ F++++E+L +v++ ++ e+e GL k+ + +A+
50365 464 QAQRKQIDRVIALFQLTREQLVDVQAKMRAELEYGLK--KKSHGLAT 508

v+kMlPtFvstPtGtEkGdFLALDLGGTNfRVllVklgngkgvemtqs
v+MlPt+V+++P+GtEkG FLALDLGGTNfRVllVlK++ ++ +v m ++
50365 509 -VRMLPTYVCGLPDGTEKGKFLALDLGGTNfRVllVlKIRSGRRSVRMYNK 557

kYriPeelmtgenvtgeqLFDfiAecikdFmdeqfpkgkkepLpLGFTFS
++iP e+m+g tge+LFD+i++ci+dF+d+++kg + LpLGFTFS
50365 558 IFAIPLLEIMQG---TGEELFDHIVQCIADFLDYMGLKGAS--LPLGFTFS 602

FpcSQtsInegiLiRwTKGfkiGRAtnsgvEGhDVVqLLreAlkrrGafp
FPC+Q sI++g+Li WTKGfK+ ++EG+DVV +LreAlkrr +f+
50365 603 FPCRQMSIDKGTLLIGWTKGfKA-----TDCEGEDVVDMLREAIKRRNEFD 647

idVVAvvNDTVGTlmscaYtkGRGdpecetviGLIvGTgtNaCYmEemrn
+d+VAvvNDTVGT+m+c+Y+ dp+ce iGLI GTG+N CYmE mrrn
50365 648 LDIVAVVNDTVGTMMTCGYE-----DPNCE--IGLIAGTGSNMCYMEDMRN 691

Figure 2C

5/5

Hexokinase Domain 2 Alignment (cont'd)

```

IekleGkLkddIpdegrMcINmEWGaFGDngldlprTkYDvviDeeSpN
Ie++eg      eg McIN+EWG FGDng++d++rT+YD ++De+S+N
50365 692 IEMVEG-----GEGKMCINTEWGGFGDNGCIDDIRTRYDTEVDEGSLN 734

PGqQlFEKMISGmYLGEivRlilldltkeglLFkgqdsplktrgsfeTs
PG+Q++EKM+SGmYLGEivR il+dltk+gllF+gg+s++L+trg+feT+
50365 735 PGKQRYEKMTSGmYLGEIVRQIILDLTKQGLLFRGQISERLRTRGIFETK 784

vLSrIEsDpsenledvrailqtaLgletTdeerklvrrvCeaVstRAARL
+LS+IESD +++l +vr iLq+ Lgl++T+e++++v++VC avs+RAA+L
50365 785 FLSQIESD-RLALLQVRRILQQ-LGLDSTCEDSI VVKEVCGAVSRRAAQL 832

caagLAAILkirengrerlkvtVGvDGSVYklyPgFkerlaealrdll
c+agLAai++k+re g+e+l++tVGvDG++Ykl+P+F+++l+e++++l+
50365 833 CGAGLAAlVEKRREDQGLEHLRITVGVDGTLYKLHPHFSRILQETVKELA 882

pdcegseedkkvsiipAEDGSGkGAAlvaAVaakl<-* (SEQ ID NO:4)
p+c      d v++ +EDGSGkGAAl++AVA +l
50365 883 PRC-----D--VTFMLSEdGSGKGAALITAVAKRL 910 (SEQ ID NO:2)

```

Figure 2D

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 November 2001 (29.11.2001)

PCT

(10) International Publication Number
WO 01/090325 A3

(51) International Patent Classification⁷: C12N 9/00, 9/04

(21) International Application Number: PCT/US01/16549

(22) International Filing Date: 21 May 2001 (21.05.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/205,508 19 May 2000 (19.05.2000) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 60/205,508 (CIP)
Filed on 19 May 2000 (19.05.2000)

(71) Applicant (for all designated States except US): MIL-LENNIUM PHARMACEUTICALS, INC. [US/US]; 75 Sidney Street, Cambridge, MA 02139 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MEYERS, Rachel, E. [US/US]; 115 Devonshire Road, Newton, MA 02468 (US). WILLIAMSON, Mark [US/US]; 15 Stonecrest Drive, Saugus, MA 01906 (US).

(74) Agents: MYERS, Louis et al.; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
18 July 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 50365, A HEXOKINASE FAMILY MEMBER AND USES THEREOF

(57) Abstract: The invention provides isolated nucleic acids molecules, designated 50365 nucleic acid molecules, which encode novel hexokinase members. The invention also provides antisense nucleic acid molecules, recombinant expression vectors containing 50365 nucleic acid molecules, host cells into which the expression vectors have been introduced, and nonhuman transgenic animals in which a 50365 gene has been introduced or disrupted. The invention still further provides isolated 50365 proteins, fusion proteins, antigenic peptides and anti-50365 antibodies. Diagnostic methods utilizing compositions of the invention are also provided.

WO 01/090325 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/16549

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 C12N9/00 C12N9/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EMBL, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL 'Online! AI949849, 14 March 2000 (2000-03-14) "National cancer Institute, Cancer Genome Anatomy Project, Tumor Gene Index http://www.ncbi.nlm.nih.gov/ncicgap " XP002191716 99% identity in 514 nt overlap (pos. 3653-3140:1- 514) with Seq. ID No. 1. ---	1-29
P, X	DATABASE EMBL 'Online! AK026414, 29 September 2000 (2000-09-29) SUGANO S ET AL: "NEDO human cDNA sequencing project" XP002191717 99,8 % identity in 1527 nt overlap (pos. 586-2112:1-1525) with Seq. ID No.3 --- -/--	1-29

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

28 February 2002

Date of mailing of the international search report

20/03/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
 Fax: (+31-70) 340-3016

Authorized officer

GONCALVES M L F C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/16549

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>DATABASE EMBL 'Online! BE159085, 2 July 2000 (2000-07-02) DIAS NETO E ET AL: "Shotgun sequencing of the human transcriptome with ORF expressed sequence tags" XP002191718 98,5 % identity in 469 nt overlap (1749-1281:18-486) with Seq. ID No. 3</p>	1-29
X	<p>DATABASE EMBL 'Online! AW475040, 2 March 2000 (2000-03-02) NCI-CGAP: ""National Cancer Institute, Cancer genome Anatomy Project, Tumor Gene Index http://www.ncbi.nlm.nih.gov/ncicgap" XP002191719 100% identity in 601 nt overlap (2524-1924:1-601) with Seq. ID No. 3</p>	1-29
A	<p>MATHUPALA S P ET AL: "ARREST OF PROLIFERATION OF HIGHLY GLYCOLYTIC TUMORS UPON TYPE II HEXOKINASE DOWN REGULATION VIA AN ANTISENSE RNA APPROACH" PROCEEDINGS OF THE ANNUAL MEETING OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH, NEW YORK, NY, US, vol. 40, March 1999 (1999-03), page 22 XP002943564 ISSN: 0197-016X abstract</p>	1-29
A	<p>KATABI M M ET AL: "HEXOKINASE TYPE II: A NOVEL TUMOR-SPECIFIC PROMOTOR FOR GENE-TARGETED THERAPY DIFFERENTIALLY EXPRESSED AND REGULATED IN HUMAN CANCER CELLS" HUMAN GENE THERAPY, XX, XX, vol. 10, 20 January 1999 (1999-01-20), pages 155-164, XP002936210 ISSN: 1043-0342 abstract</p>	1-29
A	<p>SAMIR S D ET AL: "HUMAN HEXOKINASE II: SEQUENCE AND HOMOLGY TO OTHER HEXOKINASES" BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, ACADEMIC PRESS INC. ORLANDO, FL, US, vol. 197, no. 1, 30 November 1993 (1993-11-30), pages 68-74, XP002014194 ISSN: 0006-291X abstract</p>	1-29